

Mercury and Methylmercury in the children hair and fish mostly consumed in Cubatão, São Paulo State, Brazil

Mercúrio e Metilmercúrio em cabelos de crianças e em peixes mais consumidos em Cubatão, estado de São Paulo, Brasil

RIALA6/1601

Luciana Aparecida FARIAS¹, Deborah Inês Teixeira FÁVARO^{2*}, Paulo Tadeu Meira Silva OLIVEIRA², Elisabete de Santis BRAGA³

*Correspondence address: ²Laboratório de Análise por Ativação Neutrônica, Instituto de Pesquisas Energéticas e Nucleares, IPEN-CNEN/SP. Av. Professor Lineu Prestes 2242, CEP: 05508-000, São Paulo, SP, Brazil. Phone: 55 11 31339977, FAX: 55 11 31339960, e-mail: defavaro@ipen.br

¹Departamento de Ciências Exatas e da Terra, Universidade Federal de São Paulo, UNIFESP

³Laboratório de Nutrientes, micronutrientes e traços nos oceanos - LABNUT, Instituto Oceanográfico, Universidade de São Paulo
Research sponsored by Brazilian Research Agencies FAPESP (process n. 2005/50769-2 and 2009/18233-6) and CNPq, (process n.304883/2006 -9)

Recebido: 21.03.2014 - Aceito para publicação: 30.06.2014

ABSTRACT

This study evaluated Hg and MeHg concentrations in fish muscle and children hair from Cubatão, using as strategies: i) Hg and MeHg determinations in the mostly consumed fish species, ii) Hg and MeHg determinations in 93 children hair samples (from 04 to 14 years old), and iii) analysis on the children diet consumption questionnaire answered by the respective parents. The results obtained in a similar study in Cananéia were used as references. These two regions show different environmental anthropogenic impacts and distinct eating habits. The quantity and the frequency of fish consumption were the most important factors of Hg bioaccumulation in children hair. In both cities, 95 % of mercury contents in hair were $>1 \text{ mg.kg}^{-1}$ (US EPA reference) in analyzed children, and MeHg were between 67 and 83 % of total Hg. The median for total Hg concentration in fish muscle (wet basis) ranged from 10 to $179 \text{ } \mu\text{g.kg}^{-1}$ for Cananéia and 10 to $181 \text{ } \mu\text{g.kg}^{-1}$ for Cubatão, values lower than the Brazilian limits for Hg in fish. Considering the Hg contents detected in the mostly consumed fish and hair, they indicate that the children are not at risk for Hg and MeHg contamination.

Keywords. mercury, methylmercury, children, hair, fish, food habit.

RESUMO

Este estudo avaliou a concentração de Hg e MeHg em amostras de peixes e de cabelos de crianças de Cubatão, usando-se como estratégia: i) Hg total e MeHg em peixes mais consumidos; ii) Hg total e MeHg em cabelos de 93 crianças (04 a 14 anos); iii) análise do inquérito alimentar respondido pelos pais. Os dados obtidos em estudo similar realizado em Cananéia foram usados como referência. Essas duas regiões estuarinas, sob diferentes graus de impacto ambiental, apresentam diferentes hábitos alimentares. A quantidade e a frequência de consumo de peixes foram os fatores mais importantes na bioacumulação de Hg nos cabelos. Em ambas as cidades, 95 % das concentrações de Hg total nos cabelos foram abaixo de 1 mg.kg^{-1} (limite US EPA) e o percentual de MeHg ocorreu entre 67 e 83 % em relação ao teor de Hg total. As medianas para Hg total nos músculos (base úmida) variaram de 10 a $179 \text{ } \mu\text{g.kg}^{-1}$ para peixes de Cananéia e 10 a $181 \text{ } \mu\text{g.kg}^{-1}$ de Cubatão, inferiores ao limite da legislação brasileira para Hg. A partir dos teores para Hg total e MeHg em peixes mais consumidos e cabelos, as crianças parecem não estar em risco de contaminação.

Palavras-chave. mercúrio, metilmercúrio, cabelo, crianças, peixe, hábito alimentar.

INTRODUCTION

Recent studies of European and Asian fish consumers revealed significant associations between high fish consumption and high mercury (Hg) and methylmercury (MeHg) levels^{1,2}. Fish consumption is considered the primary pathway for human accumulation of Hg and MeHg. This, in turn, results in statistically significant differences between high and low fish consumer groups³. In spite of the possible negative effects upon human health, fish consumption remains a very important protein source in the human diet. Fish is also a primary source of nutrients for child development. Many articles^{4,5} and a review⁶ discussing the risk-benefit considerations of fish consumption have been published in recent years. In the Mozaffarian and Rimm⁴ study, the conclusion presented was that for major health outcomes among adults, based on both the strength of the evidence and the potential magnitudes of the effects, the benefits of fish intake outweigh the potential risks. According to Mahaffey et al study⁵, essential polyunsaturated fatty acids, such as docosahexaenoic acid (DHA), confer benefits to brain and visual system development in infants and reduce risks of certain forms of heart disease in adults which mitigate against CH₃Hg effects. For women of childbearing age, benefits of modest fish intake, with the exception of a few selected species also outweigh risks. Myers et al⁷ presented a review about nutrient and MeHg exposure from consuming fish giving emphasis to the findings from a large prospective cohort study of a population that consumes fish daily, the Seychelles Child Development Study. Preliminary findings from this study suggest that the beneficial influence of nutrients from fish may counter any adverse effects of MeHg on the development nervous system. Facing this dilemma of risk and benefit of fish consumption FAO and WHO held an Expert Consultation Workshop in 2011⁸. Some conclusions from this workshop were: consumption of fish provides energy, protein and a range of essential nutrients; eating fish is part of the cultural traditions of many peoples and in some populations, fish is a major source of food and essential nutrients; and among infants, young children and adolescents, evidence is insufficient to derive a quantitative framework of health risks and benefits. However, healthy dietary patterns that include fish consumption and are established early in life influence dietary habits and health during adult life. Some recommendations were also described such

as develop, maintain and improve existing databases on specific nutrients and contaminants, particularly methylmercury and dioxins, in fish consumed in their region and develop and evaluate risk management and communication strategies that both minimize risks and maximize benefits from eating fish⁸.

Generally in Brazil, the communities such as indigenous groups, riverside populations and coastal dwellers (low-income groups) mainly obtain protein from fish consumption due to the fact of cheap and higher supply and cultural fishing habits. On the other hand, high-income communities have more protein options and for this reason do not frequently consume large amounts of fish. A similar trend was observed in the study conducted by Farias et al⁹ that studied children from 3 public schools (PS₁, PS₂ and PS₃) in Cananéia city observing a direct relation between total Hg content in hair and the low economic level of the family. In general, most families of the 3 schools studied presented lower income (1 to 5 minimum salaries, US\$ 300 to 1,500/month). These low-income families showed a greater dependency of protein intake from aquatic origin, in relation to bovine protein due to a greater availability and lower cost in the coastal region⁹. The majority of the studied families stated that fish consumption is derived from the families' fishing activities. Garcia¹⁰ in his study also observed the same relationship.

Continuing the study started in Cananéia, similar study was developed in Cubatão, considering the importance of assessing mercury levels in hair and commercial fish in coastal populations and their association with eating habits¹¹.

The purpose of this study was to evaluate total Hg and MeHg concentration in fish muscle and children hair, using as strategy: *i*) total Hg and MeHg determinations in the most consumed fish species; *ii*) total Hg and MeHg determination in 93 children hair samples (aged 4 to 14) in Cubatão and *iii*) analyses of answers to a diet consumption questionnaire by the parents of the children participating in the study. The results obtained for Hg in children hair in a previous study in Cananéia⁹ were used for comparison and discussion purposes. These two estuarine regions are undergoing different environmental anthropogenic impacts¹²⁻¹⁵. Cananéia is considered a naturally preserved area and recognized as a World Biosphere Reserve by UNESCO. Cubatão is located in an area under strong environmental impact caused by complex industrial activities. Eating habits

of these local populations are also undergoing different socio-economic changes. From the results of this study it is hoped that an estimation of the children population exposure to Hg and MeHg by means of hair analysis and fish intake can be obtained. Since, there are no references values for total Hg in children's hair, this study can also be used as a contribution for a pool of referential values in hair for Brazilian coastal children populations. In Brazil, total Hg levels in children hair have mostly been focused on different Amazonian communities^{16,17}.

MATERIAL AND METHODS

Studied areas

The Baixada Santista is a coastal area located in São Paulo State that includes four important coastal cities: São Vicente, Santos, Guarujá and Cubatão. Santos city has the busiest South American port. Cubatão city is one of the country's most important industrial poles with a population of 118,720 inhabitants¹⁸. Santos Bay, Santos and São Vicente estuarine channels and rivers are part of an important hydrological polluted system, which includes Cubatão city's industrial center and complex with over a hundred industries including the largest Brazilian steel plant¹⁹. Nowadays, there has been a reduction of hydrological pollution but not enough to reestablish the base-line values. In regards to Hg, some studies offer some environmental data for this region²⁰⁻²² revealing Hg contamination in sediments.

Cananéia city is located in the south coast of São Paulo state, far from the Santos-Cubatão system and is a part of the estuarine-lagoon complex of Cananéia-Iguape. In the southern part, Cananéia is surrounded by a preserved estuarine system^{15,23} that includes the Comprida, Cananéia and Cardoso Islands. This integral system is recognized by UNESCO (United Nations Educational, Scientific and Cultural Organization) as a World Biosphere Reserve. Cananéia city is also one of the oldest cities in Brazil and has one of the largest preserved natural Brazilian mangrove areas. The city has two main economic activities, which are tourism, and fishing and its 12,226 inhabitants make their living primarily of these two activities¹⁸. The extensive mangrove provides a rich breeding and feeding ground for an extremely wide variety of aquatic fauna and flora^{10,23}. This scenario contributes to the presence of a large fishing population, which consumes large quantities of fish and seafood. Figure 1 presents the regions studied.




Figure 1. Map from Cananéia and Cubatão cities, Brazil

Sample Collection

Hair sampling and preparation

The hair samples from the children (4-14 years old) from the two cities were collected at the schools indicated by the Cities Educational Departments observing that sampling was to be in communities close to the estuarine system and/or where the students were members of a fishing family. In Cubatão, hair (93) and fish (58 individuals) samples were collected in March 2008 and in Cananéia⁹, 105 hair samples and 53 fish samples were obtained in August 2007. The most adequate communities to participate in this study were located at: i) Jardim Casqueiro and Ilha Caraguatá in Cubatão and ii) Vila Cabana, Acarau and Carijo in Cananéia. These communities were also chosen based on the number of fishing families located in each region. Children from public schools (PS) in both cities were invited to participate in the study. All parents and/or guardians signed a consent agreement for their children's participation in the study. Excluding factors were those children whose parents did not sign the consent agreement and children whose hair was too short to cut, mainly boys. In order to ascertain the daily dietary intake a questionnaire was filled out by the participating families about fish weekly intake, fish species most consumed, age, gender, social and economic status. This kind of questionnaire was used in a previous study⁹. The children selected had to present the signed consent agreement, have enough hair for collection and also a filled out questionnaire. Children hair samples were collected with a single cut from the

occipital region very close to the scalp with steel scissors, in the approximate amount of 1g. A small recipient containing 1 g of hair sample was used for comparison during *in situ* sampling. The study was approved by the Ethics Committee of the Nuclear and Energy Research Institute (IPEN/CNEN-SP) (Project n^o 109/CEP-IPEN/CNEN-SP)⁹.

Fish selection and preparation for analysis

Tabulated data from this study identified the 5 most consumed fish species in Cubatão which were: three carnivorous species: *Macrodon ancylodon* (King weakfish – Pescada*), *Menticirrhus americanus* (Southern king croaker - Perna de Moça*) and *Micropogonias furnieri* (Whitemouth croaker – Corvina*) and two planktivorous species, *Mugil liza* (Mullet – Tainha*) and *Sardella braziliensis* species (Sardine – Sardinha*). In Cananéia only four fish species were cited on the questionnaires which were: three carnivorous species, *Macrodon ancylodon* (King weakfish – Pescada*), *Centropomus parallelus* (Fat snook – Robalo peba*) and *Micropogonias furnieri* (Whitemouth croaker – Corvina*) and one planktivorous species, *Mugil liza* (Mullet – Tainha*). The families that took part in this study informed that they usually bought fish from the local markets, except the fishing families. Due to this, all the fish samples analyzed in this study were bought in local markets as well. (*Popular fish name in Portuguese).

The fish samples were bought directly from local street markets in Cubatão during March of 2008 and in Cananéia city in August 2007. All samples were assessed for quality indicators and external organoleptic characteristics (eyes, gills and scales). Fish were conditioned in isothermic boxes in crushed ice (-4 °C) and then identified before registering the biometric information (total length, total weight and body weight). In the laboratory, the samples of fish muscle were obtained from the 58 fishes from Cubatão and the 53 from Cananéia. The samples were dried at 40 °C in a ventilated oven until constant weight. Dried samples were ground, homogenized and prepared for total Hg determination. These dried samples were then analyzed and consequently, the results for total Hg and MeHg were obtained on a dry weight basis. After that, dry weight results obtained were transformed to wet weight basis by using humidity %.

Total Hg and MeHg determination

Total Mercury and MeHg determinations

were performed by Cold Vapor Atomic Absorption Spectrometry (CV AAS), using a FIMS (Flow Injection Mercury System) from Perkin Elmer. The experimental procedure applied in hair and fish samples have already been described in previous papers^{9,24}. The methodology validation for total Hg and MeHg determinations was carried out using human hair reference material (IAEA-086), Dogfish muscle (DORM-1, NRCC) and Dogfish liver (DOLT-3, NRCC), for fish samples. The detection limit (DL) for total Hg determination is 0.001 mg.kg⁻¹ and quantification limit (QL), 0.010 mg.kg⁻¹ 24.

All analyses were performed at the Neutron Activation Analysis Laboratory in Nuclear and Energy Research Institute – LAN/IPEN, SP-Brazil.

Statistical treatment

The statistical analysis was processed by the IBM SPSS Statistics Version 17.0 software (Tulsa, OK, USA).

For fish and total Hg in children hair from the different public schools investigated, the Shapiro Wilkis data treatment was used. However, since no normal Hg level distribution was found, the non-parametric Kruskal Wallis was applied to verify the significance of the differences in the Hg levels in the fish samples. The Spearman correlation test was used for the Hg and other variables.

RESULTS AND DISCUSSION

The precision and accuracy for total Hg and MeHg determinations were checked by means of reference material analyses in fish muscle (Dorm-1 – Dogfish muscle and Dolt-3 – Dogfish liver) and hair (IAEA-086). The results for total Hg and MeHg determinations in the IAEA-086 reference material showed relative standard deviation (RSD) of 3.6 and 6.3 % (n=10) and relative error (RE) of 1.0 and 1.5 %, respectively. For fish samples, the results showed RSD from 4.2 to 11.7 and RE from 0.9 to 12.0% (n=10). These results showed good precision and accuracy for both analytical methodologies in both matrices.

The fish samples were submitted to a biometric procedure that was applied to each individual before the muscle sampling and mercury analyses. Table 1 presents the total body length, feeding habits, median and interval for total Hg and MeHg (%) concentrations in fish muscle of the different species analyzed from Cananéia and Cubatão cities. In general, fish from a non-polluted costal region (without effective Hg contamination) present total Hg concentrations around

0.15 mg.kg⁻¹ (wet weight) whereas fish in contaminated regions can reach total Hg levels up to 2 mg.kg⁻¹ (wet weight). Benthic species can reach concentrations from 10 to 20 mg.kg⁻¹, values considered lethal for fish²⁵. In the present study, the higher total Hg levels found in Cananéia and Cubatão fishes, were 0.442 mg.kg⁻¹ and 0.348 mg.kg⁻¹ (wet weight), respectively, both in *Micropogonias furnieri* (Whitemouth croaker), a detritivorous species. These values are lower than the maximum levels established by the current Brazilian legislation (2013)²⁶ considering predatory and non-predatory fishes, 1.0 and 0.5 mg.kg⁻¹, respectively. In the present study, total Hg values for the individuals analyzed ranged from < 0.010 to 0.442 mg.kg⁻¹ (median of 0.010 mg.kg⁻¹ for *Mugil liza* (Mullet) to 0.179 mg.kg⁻¹ for *Micropogonias furnieri* (Whitemouth croaker), for fish available for consumption in Cananéia city. For

Cubatão, the values ranged from < 0.010 to 0.348 mg.kg⁻¹ (median of 0.010 mg.kg⁻¹ for *Mugil liza* to 0.181 mg.kg⁻¹ for *Micropogonias furnieri*). All data are presented in wet weight. According to the limits proposed for fish and fish products, considering the predatory and non-predatory species sampled, none surpassed the limits established by Brazilian legislation²⁶.

The percentage of MeHg determined in fish muscle showed values ranging from 14 % to 99 % taking into account the total Hg level. In general, the predominant form of Hg in fish muscle is methylmercury²⁷. In other tissues such as liver and kidney Hg presents itself as inorganic form exceeding the values of Hg found in the muscle^{25,28}. An exception is the Pacific blue Marlin that presents a 20 % level of MeHg in its muscle tissue. In general, the % of MeHg follows the pattern: muscle >

Table 1. Total length, weight, feeding habits, median and range for total Hg and MeHg (%) content in different fish species analyzed from Cananéia and Cubatão

Fish species (n)*	Feeding Habits	Weight (g)	Total length (mm)	Total Hg (mg.kg ⁻¹) (wet weight)	Total Hg (mg.kg ⁻¹) (dry weight)	% MeHg (interval)
					Muscle	Muscle
CANANÉIA						
Whitemouth croaker (11)	Detritivorous	1134 832 - 1711	454 414 - 507	0.179 0.114-0.442	0.812 0.516 - 2.008	63 (34 - 85)
King weakfish (16)	Carnivorous	298 220 - 613	320.5 200 - 385	0.039 0.012- 0.100	0.180 0.056 - 0.456	66 (37 - 99)
Fat snook (12)	Carnivorous	481 312 - 584	399.5 318 - 424	0.032 0.015-0.178	0.129 0.061 - 0.712	45 (18 - 67)
Mullet (14)	Planctivorous	1514 807 - 1936	567.5 432 - 600	0.010 <0.010-0.025	0.032 <0.010 - 0.098	---
CUBATÃO						
Whitemouth croaker (11)	Detritivorous	720 607 - 909	426 360 - 453	0.181 0.041-0.348	0.789 0.178 - 1.512	37 (18 - 61)
Southern king croaker (12)	Carnivorous	1736 1446 - 2170	324 301 - 385	0.105 0.043-0.184	0.499 0.206 - 0.878	53 (22 - 68)
King weakfish (16)	Carnivorous	304 234 - 374	355 200 - 368	0.046 0.012-0.062	0.193 0.051-0.257	42 (18 - 61)
Sardine (16)	Planctivorous	104 91 - 120	223 210 - 235	0.027 <0.010-0.040	0.091 0.026 - 0.135	31 (14 - 61)
Mullet (5)	Planctivorous	1416 1302 - 1710	547 520 - 555	0.010 <0.010-0.023	0.029 0.013 - 0.090	56 (56 - 68)

n: number of samples analyzed; --- : Values for MeHg < 0.010 mg.kg⁻¹

gills > kidneys > liver²⁹. Some studies verified that the % of MeHg in muscle tissues is higher in upper trophic levels. In marine ecosystems the % of MeHg was higher in tertiary carnivorous species (C₃) than secondary ones (C₂) while their total Hg levels were of the same order. These observations demonstrated the biomagnification capacity of Hg throughout the trophic food chain and reflect the fact that the upper trophic fish present higher levels of MeHg³⁰.

Observing the data assembly according to Kruskal-Wallis statistical treatment, the differences between Total Hg observed in function of the fish species analyzed (*carnivorous and non-carnivorous*) presented a high significant difference ($p < 0.05$), as expected, once carnivorous species are exposed to a more intense Hg bioaccumulation rate than the non-carnivorous species³⁰




Figure 2. Total Hg concentrations distribution in hair samples in female and male groups from Cananéia and Cubatão cities, Brazil

Table 2. Total Hg and % MeHg results in children's hair from Cubatão public schools and weekly fish intake

(Total Hg – mg.kg ⁻¹)	CUBATÃO		CANANÉIA ^a		
	PS _{Ct1}	PS _{Ct2}	PS _{Ca1}	PS _{Ca2}	PS _{Ca3}
Female (n)	31	36	7	26	35
Mean ± std	0.236 ± 0.145	0.338 ± 0.345	0.041 ± 0.035	0.462 ± 0.258	0.461 ± 0.361
Median	0.235	0.324	0.033	0.394	0.326
range	0.016 – 0.637	< 0.010 – 1.330	< 0.010 – 0.112	< 0.010 – 0.959	0.038 – 1.479
Age (median) (range)	11 (9-14)	10 (7-13)	5 (4-9)	8 (5-12)	6 (4-12)
% MeHg (median)	74	71	74	69	79
Male (n)	13	13	8	12	17
Mean ± std	0.458 ± 0.169	0.483 ± 0.554	0.249 ± 0.262	0.636 ± 0.881	0.745 ± 0.787
Median	0.467	0.296	0.203	0.379	0.506
range	0.220 – 0.741	<0.010 – 1.859	< 0.010 – 0.770	0.076 – 3.321	< 0.010 – 2.814
Age (median) (range)	10 (9-12)	10 (8-12)	6 (4-8)	9 (3-10)	7 (5-10)
% MeHg (median)	83	72	68	83	83
Total (n)	44	49	15	38	52
Mean ± std	0.301 ± 0.182	0.376 ± 0.362	0.152 ± 0.216	0.641 ± 0.537	0.576 ± 0.547
Median	0.270	0.319	0.043	0.384	0.405
Range	0.016 – 0.741	< 0.010 -1.859	<0.010 – 0.770	<0.010 – 3.321	< 0.010 – 2.814
age (median) (range)	11 (9-14)	10 (7-13)	5 (4-9)	8 (4-12)	7 (4-12)
% MeHg (median)	74	71	72	70	81
Fish weekly intake (%)					
Rarely	34	44	13	8	7
1 - 2 times/week	57	49	48	45	43
> 2 times/week	9	7	39	47	50

a: Farias et al⁹; n: number of individuals; PS_{Ct1, Ct2}: are public schools from Cubatão; PS_{Ca1, Ca2, Ca3}: are public schools from Cananéia

in function of its position in the food chain.

Table 2 shows the mean, median and interval for the total Hg results in children's hair, as well as, median age, MeHg (%) and fish weekly intake for female and male groups, from Cananéia and Cubatão communities. Some data for total Hg in children hair from Cananéia city obtained in a previous study of our group were already published⁹. Hair mercury content was lower than the US EPA reference limit³¹ of 1 mg.kg⁻¹ in 95 % of the studied children; 4 % presented levels between 1 and 2 mg.kg⁻¹ and 1 %, higher than 2 mg.kg⁻¹. The normal level of Hg in hair is 1-2 mg.kg⁻¹, however those who consume fish one or more times per day may have Hg levels in hair exceeding 10 mg.kg⁻¹³².

Figure 2 shows the total Hg concentrations distribution in hair samples in female and male groups from Cananéia and Cubatão cities. The mean values for male groups were higher in both counties (Table 2). Nakagawa (1995), *apud* Yasutake et al³³ observed that gender can be an important variable for total Hg concentration in hair and verified that the content in males is much higher than in females. On the other hand, there are studies where a significant correlation between gender and total Hg content in hair was not observed³⁴. Presently, there is still no agreement about the influence of gender on the total Hg content in hair.

In the present study, a negative correlation between age and total Hg among the entire sampled group was also found. The Spearman correlation coefficient was found to be $r_s = -0.336$, indicating a weak negative correlation between age and total Hg content in children's hair.

The PS_{Ct1} School individuals sampled from the Cubatão region presented a total Hg mean of 0.236 mg.kg⁻¹ for females and 0.458 mg.kg⁻¹ for the male group. At the PS_{Ct2} School, the mean was 0.338 mg.kg⁻¹ for females and 0.483 mg.kg⁻¹ for the male group. Cananéia region, at the PS_{Ca1} School, individuals presented total Hg mean of 0.041 mg.kg⁻¹ for the female group, 0.249 mg.kg⁻¹ for the male group; PS_{Ca2} School, 0.462 mg.kg⁻¹ for the female group, 0.636 mg.kg⁻¹ for the male group and PS_{Ca3} School, 0.461 mg.kg⁻¹ for the female group and 0.745 mg.kg⁻¹ for the male group. Table 2 also shows the median of MeHg (%) in the hair samples analyzed. The MeHg % ranged from 71 to 83 % for both groups in Cubatão and from 68 to 83 %, in Cananéia. In the literature, the average value of 80 % of MeHg in relation to total Hg content in hair is commonly found³⁵.

According to the questionnaire answers regarding

eating habits in Cananéia city (Table 2), a frequency of 1-2 times/week of fish and seafood intake occurred in 48 %, 45 % and 43 % for children from PS_{Ca1}, PS_{Ca2} and PS_{Ca3}, respectively. Furthermore, according to the questionnaire data 39, 47 and 50 % reported consuming fish > 2 times/week, respectively. Rare fish consumption was reported by 13 %, 8 % and 7 %, respectively, showing the importance of fish in the diet of the local population. In general, the sampling revealed that most families (67.6 %) belonged to a low socio-economic level, receiving 1 to 5 minimum salaries in Brazil (US\$ 300 to 1500), contributing to a higher fish and seafood consumption due to the fact that most families make their living by fishing⁹.

In Cananéia, the children hair samples from public school PS_{Ca1} presented median values for total Hg, for both male and female groups, much lower than the PS_{Ca2} and PS_{Ca3} values. Furthermore, schools PS_{Ca2} and PS_{Ca3} presented the higher number of fishermen as parents (45 %). This fact may explain why these children consume more fish/seafood and as a result presented higher total Hg levels in their hair than the children from PS_{Ca1}.

In Cubatão, the children hair samples from public schools PS_{Ct1} and PS_{Ct2} presented very similar median values for total Hg (0.270 and 0.319 mg.kg⁻¹) and higher mean values for the male group in both schools as well (Table 2).

According to eating habits in Cubatão city, the present study verified, observing the parents answers to the questionnaire (Table 2), that a 1-2 times/week of fish and seafood intake frequency occurred in 57 % and 49 % for PS_{Ct1} and PS_{Ct2}, respectively. Furthermore, 9 and 7 % reported consuming fish more than 2 times/week. Rare fish consumption was observed in 34 % and 44 %, respectively. These results indicate a different frequency in fish and seafood consumption in Cubatão when compared to the fish consumption data from Cananéia. This difference may also explain the higher mean Hg levels in children's hair (female and male groups) in Cananéia (except for PS_{Ca1}) than Cubatão as Hg concentration in fish species most consumed were similar in both communities. This seems to confirm the influence of higher fish intake frequency in Cananéia.

In Cubatão, the sampling also revealed that most families (59 %) belonged to a low socio-economic level, receiving 1 to 5 minimum salaries. The majority of the parents interviewed answered that they consumed fish acquired in the weekly open street market.

Table 3. Total Hg concentrations in children's hair from different localities in Brazil and other countries

Brazil	Total Hg (mg kg ⁻¹)	
	Mean ± s.d.	Interval
São Luiz do Tapajós – age group: 0 - 5 years (Pará State – Northern Brazil) ²⁵	21.06 ± 14.38	0.10 – 94.5
Barão do Melgaço – age group: 3 – 7 years (Mato Grosso State – Central Brazil) ³⁸	5.37 ± 3.35	0.58 – 17.14
River Madeira basin: children and adults (Amazon region – Northern Brazil) ¹⁷	15.2 ± 9.6	0.36 – 150
Manaus city – age group: 2 – 7 years (Amazon region- Northern Brazil) ¹⁶	1.93 ± 3.90	0.02 – 34.4
Cubatão – age group: 1- 10 years (São Paulo State - Southeastern Brazil) ¹⁹	0.88 ± 0.61	–
Cananéia – age group: 4 - 12 years (São Paulo State- Southern Brazil) ⁹	0.46 ± 0.35	< 0.01 – 3.33
Cubatão – age group: 4 - 14 anos (São Paulo State - Southern Brazil) (present study)	0.34 ± 0.29	< 0.01 – 1.86
Other countries	Total Hg (mg kg ⁻¹)	
	Mean ± s.d.	Interval
Spain – age: 4 years ²	0.99 ± 0.20	< 0.01 – 5.63
Japan – age: 7 years ³³	1.4 ± 0.89	0.45 – 6.32

The available information in regards to the Hg content in children hair from different Brazilian regions is focused primarily on data from the Amazon region^{16,17}. Hacon et al³⁶ and Passos and Mergler³⁷ published two reviews showing the mercury concentration in the Amazon.

Table 3 presents total Hg levels in children hair from Cananéia⁹ and Cubatão (present study) and children living in the Amazon region and two other countries. Total Hg levels in children hair found in the present study were much lower when compared to values obtained from the Amazon region and lower than values published by ACPO¹⁹. The values observed in this study were also lower than values from other countries, despite the difficulty for comparison based on age differences between participants. However, considering these different studies, fish consumption is often cited as the most significant factor for human mercury bioaccumulation, particularly in its organic form. The results presented in this study suggest that the concentration of total Hg and MeHg found

in children hair samples were mainly dependent on eating habits of the population, the same observed by Tavares et al³⁸ and Marques et al³⁹ in studies developed with children in Brazil as well as low economic level families.

It was expected that the highest Hg and MeHg levels were to be found in children hair living in Cubatão in relation to those observed in children living in the Cananéia city due to the association of the fish source from the polluted estuarine system in Cubatão (industrial activities and the largest Brazilian steel plant) as mentioned before. However, this was not the case. The fish consumed in Cubatão did not necessarily come from local waters. Furthermore, there are fewer fishing families in this industrial region. On the other hand, the number of fishing families in Cananéia city was much higher than in Cubatão, probably resulting in the higher fish consumption in Cananéia. In both locations, the families were classified as low-income varying from 1 to 5 minimum salaries (67.6 % in Cananéia and 59 % in Cubatão) according to POF (Family Budget Survey -

POF 2008-2009 (the Brazilian census¹⁸). This fact also seems to favor fish/seafood consumption for protein in coastal cities. It should be noted that Cubatão city offers more options for protein sources than Cananéia. This behavior could be associated to the high price of fish and to the pollution estuarine history of Cubatão which may contribute to lower fish consumption. On the other hand, Cananéia estuary has an extensive mangrove area that provides a rich breeding and feeding ground for an extremely wide variety of aquatic fauna and flora, producing a large amount of fish and seafood.

In relation to total Hg fish concentrations, both Cananéia and Cubatão fish presented levels below the fish Hg limits established by Brazilian legislation for human consumption (predatory and non-predatory fishes, 1.0 and 0.5 mg.kg⁻¹, respectively). In Cubatão city, the origin of the fish in the local markets was verified by questioning the sales person during the purchase. They stated that most commercialized fish originated from the coastal areas of Santa Catarina State, in the southern Brazilian region, far from this polluted region.

Researchers from Duke University, USA, recently published a study about photolytic degradation of MeHg in brackish waters where the Hg concentrations in freshwater are much greater than those found in salt water. However, when fish are ingested, it is the saltwater fish that offer greater risk to human health⁴⁰. The MeHg is potentially more dangerous than inorganic Hg forms and in the water system is associated with dissolved organic matter and in the sea, is also associated with chloride, ie, salt. One way nature transforms the MeHg in a less toxic form is through photochemical reactions promoted by sunlight. When MeHg is associated to dissolved organic matter such as plants or animals detritus, the sunlight breaks the molecule. However, in ocean water, the MeHg remains strongly linked to chloride and is not easily degraded by sunlight. Since MeHg is not broken down by sunlight in sea water, its residence time is much greater in saltwater than in freshwater, which increases the rate of intake by marine animals⁴⁰. Fish and seafood have a natural tendency to store MeHg in their bodies characterized therefore as the main source of metal intake in humans. In the marine environment, bacteria can methylate the Hg and transform it into an easily assimilated form especially in surface marine sediments. According to the same study it is also important to direct research and policy control of Hg for saltwater, mainly in the coastal areas those are submitted to anthropogenic

inputs. To date, most efforts have been directed to the presence of the metal in freshwater. This fact is particularly true in Brazil, given that most studies with Hg are located in the Amazon region.

Finally, the results obtained in the present study indicated the necessity of obtaining more data concerning Hg values for different coastal regions of Brazil in order to contribute to establishing values considered characteristic for different parts of the country and in relation to human populations. Children, in particular, need intensive and continuous studies because they are most sensitive to the deleterious effects of this element.

CONCLUSION

The fish species analyzed were the most consumed by the coastal populations studied and presented Hg levels in muscles lower than the limit established by Brazilian legislation (0.5 – non-carnivorous species and 1.0 mg.kg⁻¹ – carnivorous species, respectively) in both studied regions. The median values for total Hg levels in the fish species analyzed were very similar in both coastal cities.

The Hg concentrations in children hair were lower than the US EPA reference limit of 1 mg.kg⁻¹ in 95 % of the studied children. This value was used as reference since at present there is no value set for children. The mean values for total Hg in children's hair (female and male groups) in Cananéia were higher than in Cubatão, except for PS_{Ca1} School. From these results, it seems that the quantity and frequency of fish ingested are the main factors that contribute to the exposure and bioaccumulation of Hg, as the total Hg levels in fish were similar in both communities.

To complement information about Hg in communities outside the Amazon region it is important to determine the profile of Hg and MeHg distribution in Brazil, considering the wide diversity of ecosystems, cultural patterns and socio-economic conditions particularly considering the frequency and type of fish people consume.

The results from this study can also be used as a contribution for referential values for total Hg levels in hair for Brazilian coastal children populations.

Considering the Hg levels in those fish species most consumed and hair, children from both communities studied appear not be at risk of Hg and MeHg contamination.

ACKNOWLEDGMENTS

The authors are grateful to the Municipal Education Department from Cananéia and Cubatão cities, from São Paulo State and especially to the schools participating in the study; the Oceanographic Institute of the São Paulo University for the logistic support in Cananéia City and for financial support from the Research Agencies FAPESP (process n^o 2005/50769-2 and 2009/18233-6) and CNPq (process n^o 304883/2006 - 9).

REFERENCES

1. Endo T, Haraguchi K. High mercury levels in hair samples from residents of Taiji, a Japanese whaling town. *Mar Pollut Bull*.2010;60:743-7.
2. Freire C, Ramos R, Lopez-Espinosa MJ, Díez S, Jesús V, Ballester F, et al. Hair mercury levels, fish consumption, and cognitive development in preschool children from Granada, Spain. *Environ Res*. 2010;110(1):96-104.
3. Chien LC, Gao CS, Lin HH. Hair mercury concentration and fish consumption: risk and perceptions of risk among women of childbearing age. *Environ Res J*.2010;110(1):123-9.
4. Mozaffarian D, Rimm EB. Fish intake, contaminants, and human health: evaluating the risks and the benefits. *JAMA*.2006;296(15):1885-99.
5. Mahaffey KR, Sunderland EM, Chan HM, Choi AL, Grandjean P, Marien K, et al. Balancing the benefits of n-3 polyunsaturated fatty acids and the risks of methylmercury exposure from fish consumption. *Nutr Rev*.2011;69(9):493-508.
6. Costa, LG. Contaminants in fish: risk-benefit considerations. *Arh Hig Rada Toksikol*. 2007; 58:367-74.
7. Myers G, Davidson PW, Strain JJ. Nutrient and methylmercury exposure from consuming fish. *J Nutr*.2007;137:2805-8.
8. Food and Agriculture Organization of the United Nations and World Health Organization - FAO/WHO. Report of the Joint FAO/WHO expert consultation on the risks and benefits of fish consumption, 25-29 January 2010, Report # 979; Food and Agriculture Organization of the United Nations and World Health Organization: Rome. 2011.
9. Farias LA, Santos NR, Fávoro DIT, Braga ES. Mercúrio total em cabelo de crianças de uma população costeira, Cananéia, São Paulo, Brasil. *Cad Saúde Pública*.2008;24(10):2249-56.
10. Garcia TR. Impactos da implantação de uma cooperativa de produção de ostras junto a comunidades extrativistas caiçaras do litoral sul/SP: um estudo de caso [dissertação de Mestrado]. São Paulo (SP): Universidade de São Paulo; 2005.
11. Díez S, Montuori P, Pagan A, Sarnocchiaro P, Bayona JM, Triassu M. Hair mercury levels in an urban population from southern Italy: fish consumption as a determinant of exposure. *Environ Int*.2008;34:162-7.
12. Abessa DMS, Carr RS, Rachid BRF, Sousa ECPM, Hortelani MA, Sarkis JES. Influence of a Brazilian sewage outfall on the toxicity and contamination of adjacent sediments. *Mar Pollut Bull*.2005;50(8):875-85.
13. Amorim EP, Favaro DIT, Berbel GBB, Braga ES. Assessment of metal and trace element concentrations in the Cananéia estuary, Brazil, by neutron activation and atomic absorption techniques. *J Radioanal Nucl Chem*.2008;278 (2):485-9.
14. Braga ES, Bonetti CVDH, Burone L, Boneti Filho J. Eutrophication and bacterial pollution caused by industrial and domestic wastes at the Baixada Santista Estuarine System - Brazil. *Mar Pollut Bull*.2000;40(2):165-73.
15. Mahiques MM, Burone L, Figueira RCL, Oliveira AA, Wanderley L, Capellari B, et al. Anthropogenic influences in a lagoonal environment: A multiproxy approach at the Valo Grande Mouth - Cananéia-Iguape System (SE-Brazil). *Braz J Oceanogr*.2009;57(4):325-37.
16. Farias LA, Favaro DIT, Pessoa A, Aguiar JPL, Yuyama LKO. Mercury and methylmercury concentration assessment in children's hair from Manaus, Amazonas State, Brazil. *Acta Amazon*.2012;42(2):279-86.
17. Bastos WR, Gomes JPO, Oliveira RC, Almeida R, Nascimento EL, Bernardi JVE. Mercury in the environment and riverside population in the Madeira River Basin, Amazon, Brazil. *Sci Total Environ*.2006;368:344-51.
18. Instituto Brasileiro de Geografia e Estatística - IBGE. Pesquisa de Orçamentos Familiares 2008/2009. Análise do consumo alimentar pessoal no Brasil. [Accessed 2013 September 13th]. Available from:[http://www.ibge.gov.br/home/estatistica/populacao/condicaodevida/pof/2008_2009/default.shtm].
19. Associação de Consciência à Prevenção Ocupacional - ACPO. O Hg na Baixada Santista. [Accessed 2013 September 13th]. Available from:[<http://www.acpo.org.br/principal.php>].
20. Companhia Ambiental do Estado de São Paulo - CETESB. Sistema Estuarino de Santos São Vicente, Relatório Técnico. [Accessed 2013 September 13th]. Available from: [<http://www.cetesb.sp.gov.br/agua/aguas-superficiais/35-publicacoes/-relatorios.%202011>].
21. Hortellani MA, Sarkis JES, Bonetti J, Bonetti C. Evaluation of mercury contamination in sediments from Santos-São Vicente Estuarine System, São Paulo State, Brazil. *J Braz Chem Soc*.2005;16(6A):1140-9.
22. Luiz-Silva W, Matos RHR, Kristosch GC. Geoquímica e índice de geoacumulação de mercúrio em sedimentos de superfície do Estuário de Santos - Cubatão (SP). *Quím Nova*.2002;25(5):753-6.
23. Braga ES. Nutrientes dissolvidos e produção primária do fitoplâncton em dois sistemas costeiros do estado de São Paulo [dissertação de Mestrado]. São Paulo (SP): Universidade de São Paulo, 1995.
24. Farias LA, Fávoro DIT, Vasconcellos MBA. Determinação de mercúrio e metilmercúrio em amostras de cabelo e peixes. *Rev Inst Adolfo Lutz*.2009;68(3):451-60.
25. Malm O, Branches FJ, Akagi H, Castro MB, Pfeiffer M, Harada M, et al. Mercury and methylmercury in fish and human hair from Tapajos River Basin, Brazil. *Sci Total Environ*.1995;175(2):141-50.
26. Brasil. Ministério da Saúde. Resolução RDC n^o 42, 29 de agosto de 2013. Aprova o Regulamento técnico sobre Limites Máximos de Contaminantes Inorgânicos em Alimentos. Diário Oficial [da] República Federativa do Brasil. Brasília, DF, 30 agosto 2013. Seção 1, n^o 168, p.33-34.
27. Driscoll CT, Mason RP, Chan HM, Jacob DJ, Pirrone N. Mercury as a global pollutant: sources, pathways, and effects. *Environ Science Tech*.2013;47:4967-83.

28. Kehring HÁ, Costa M, Moreira IE, Malm O. Total mercury and methylmercury in a Brazilian estuary. *Mar Pollut Bull.*2002;44:1018-23.
29. Baêta AP. Mercúrio total e metilmercúrio em diferentes espécies de peixes da Baía de Guanabara [tese de doutorado]. Rio de Janeiro (RJ): Pontifícia Universidade Católica; 2004.
30. Gray JS. Biomagnification in marine systems: the perspective of an ecologist. *Mar Pollut Bull.*2002;45:46-52.
31. (US. Environmental Protection Agency, Office of Science and Technology, Office of Water) - USEPA, 2001. Water quality criterion for the protection of human health. Methylmercury. [Accessed 2014 March 25th]. Available from: [<http://water.epa.gov/scitech/swguidance/standards/criteria/health/methylmercury.cfm>].
32. World Health Organization - WHO, 2008. Guidance for identifying populations at risk from Hg exposure. Issued by UNEP DTIE Chemicals Branch and WHO Department of Food Safety, Zoonosis and Foodborne Diseases. [Accessed 2013 September 13th]. Available: [<http://www.who.int/foodsafety/publications/chem/mercuryexposure.pdf>].
33. Yasutake A, Matsumoto M, Yamaguchi M, Hachiya N. Current hair mercury levels in Japanese: survey in five districts. *Tohoku J Exp Medicine.*2003;199:161-9.
34. National Academy of Sciences. Toxicological Effects of Methylmercury. Committee on the Toxicological Effects of Methylmercury, Board on Environmental Studies and Toxicology, National Research Council, 2000. [Accessed 2014 March 25th]. Available on: [http://www.nap.edu/openbook.php?record_id=9899].
35. Magos L, Clarkson T. The assessment of the contribution of hair to methylmercury excretion. *Toxicol Letters.*2008;182:48-9.
36. Hacon S, Barrocas PRG, Vasconcellos ACS, Barcellos C, Wasserman JC, Campos RC, et al. An overview of mercury contamination research in the Amazon basin with an emphasis on Brazil. *Cad Saúde Pública.*2008;24(7):1479-92.
37. Passos CJS, Mergler D. Human Mercury exposure and adverse health effects in the Amazon: a review. *Cad Saúde Pública.*2008;24(Suppl 4): S503-20.
38. Tavares LMB, Câmara VM, Malm O, Santos ECO. Performance on neurological development tests by riverine children with moderate mercury exposure in Amazonia, Brazil. *Cad Saúde Pública.*2005;21(4):1160-7.
39. Marques RC, Dórea JG, Bastos R, Rebelo MF, Fonseca MF, Malm O. Maternal mercury exposure and neuro-motor development in breastfed infants from Porto Velho (Amazon), Brazil. *Int J Hyg Environ Health.*2007;210:51-60.
40. Zhang T, Hsu-Kim H. Photolytic degradation of methylmercury enhanced by binding to natural organic ligands. *Nature Geosc.*2010; 3:473-6.