

PROTEOMIC ANALYSIS OF SAFFRON (*Crocus sativus* L.) GROWN UNDER CONDITIONS OF CADMIUM TOXICITY

ANÁLISE PROTEÔMICA DO AÇAFRÃO (*Crocus sativus* L.) CULTIVADO SOB CONDIÇÕES DE TOXICIDADE POR CÁDMIO

Junfeng RAO^{1*}; Weide LV¹; Jumei YANG²

1. Hangzhou Vocational and Technical College, Hangzhou 310018, China; 2. Panan People's Hospital, Jinhua 322300, China.
raojunfenghz@163.com; 13588867756@139.com

ABSTRACT: Cd is a highly detrimental global environmental pollutant. Plants have evolved complex defense mechanisms as an adaptation to against Cd toxicity. In this study, a pot experiment was performed to evaluate the protein profile of saffron in response to Cd stress. Fifteen proteins were found to be up-regulated in the leaves of plants under Cd stress and were primarily related to metabolism, signal transduction, stress and defense response and energy. Eleven proteins were found to be down-regulated following Cd treatment, including ribulose biphosphate carboxylase/oxygenase (Rubisco), ferredoxin-NADP reductase, a 70 kDa heat shock-related protein and three protein synthesis-associated proteins. The results provide valuable insights regarding the molecular mechanism of saffron in response to Cd toxicity and the possible utilization of genetic resources in developing Cd tolerant/low-accumulation saffron.

KEYWORDS: Heavy metal. Proteome. Response. Photosynthesis. Saffron

INTRODUCTION

Cadmium (Cd) pollution has become a serious environmental issue as a consequence of human activities (KULIK et al., 2012; CAO et al. 2014a). Since the occurrence of "Itai-Itai disease" during the 1950-60s in Japan, Cd pollution has generated worldwide concern (CAO et al., 2013). Moderate Cd pollution in the soil could result in considerable Cd accumulation in cereals and subsequently pose a significant threat to human health *via* the food chain (URAGUCHI and FUJIWARA 2012). In plants, having no biological function and high toxicity, Cd can lead to various problems such as the inhibition of photosynthesis and transpiration, alteration of ion homeostasis, and poor quality products (VESELOV et al., 2003; DENG et al., 2014).

Cd affects many important physiological and biochemical processes in plants such as photosynthesis and nutrient homeostasis, which in turn inhibits normal plant growth and development (CLIJSTERS; VANASSCHE, 1985; CAO et al., 2014b). To defend against the detrimental effects of Cd, plants have evolved many detoxification mechanisms, including Cd chelation and vacuolar compartmentalization (HALL, 2002). For example, OsHMA3 (a P-type ATPase 3) has been found to play a key role in xylem Cd transport by mediating vacuolar compartmentalization in the root cells of rice (URAGUCHI; FUJIWARA, 2012). *OsDEP*, encoding a Cys-rich G protein γ subunit, has also been found to play an important role in Cd tolerance

(KUNIHIRO et al., 2013). However, the regulatory mechanism of Cd tolerance in saffron is still unknown. Saffron (*Crocus sativus*), a perennial herb belonging to the Iridaceae, is heavily used in traditional medicine because of its anticancer, antioxidative, antidepressant, anticonvulsant and learning- and memory-enhancing properties (MACCARONE et al., 2008; SHAMSA et al., 2009; ASADI et al., 2014). There is only one study which reports the effect of Cd on saffron and they only investigated the heavy metal concentration in saffron (ABOU-ARAB; ABOU DONIA 2000). However, the molecular mechanisms underlying the response to Cd toxicity in saffron remain unclear. Comparative proteomic analysis provides a powerful tool to identify the proteins expressed under biotic and abiotic stress (XU et al., 2010). No study has reported the effect of Cd on protein expression in saffron. Thus, the present study was carried out to identify the protein profiles of saffron under Cd stress. These results will be useful for understanding the effect of Cd on saffron at the proteomic level, and may identify key proteins or genes for the genetic enhancement of tolerance to Cd toxicity in plants.

MATERIAL AND METHODS

Experimental design

A pot experiment was conducted in the greenhouse of Hangzhou Vocational & Technical College, Hangzhou, China. A single saffron genotype (*Crocus sativus* L.) was used. Two kg of

commercial soil was weighed and placed in each pot (5 L, 20 cm × 22 cm × 4 cm). The soil was then artificially contaminated by adding a Cd solution to create a Cd treatment with 20 mg kg⁻¹ Cd; the control contained no added Cd. Following addition of the Cd solution, the soil was allowed to equilibrate for a period of 30 d in the greenhouse. The saffron bulbs were placed on a shaded shelf for germination. After 60 days, uniform healthy plants were selected and transplanted into the pots. Four plants were planted per pot. The experiment used a completely randomized design with 4 replicates. Plant samples were collected after 30 days of treatment for protein analysis. The fresh leaves were immediately frozen in liquid nitrogen and stored at -80 °C until analysis.

Protein analysis

Total leaf protein was extracted according to the protocol of Carpentier et al. (2005). The protein content was measured with a standard Bradford assay using bovine serum albumin as the standard (Bio-Rad, Hercules, CA, USA). Protein spots were separated using two-dimensional gel electrophoresis (2-DE) and then visualized by silver staining of the analytical gels (BAH et al., 2010; DAI et al., 2013).

The gels were scanned and calibrated with a PowerLook1100 scanner (UMAX), and protein spots were then analyzed using GE HealthCare Software (Amersham Biosciences). Only those spots with significant and reproducible changes ($P \leq 0.05$) were considered to be differentially expressed proteins. The selected protein spots were excised from the stained gels and digested with a trypsin solution using a Spot Handling Workstation (Amersham Biosciences). Trypsin digestion and peptide extraction were conducted according to the protocol of Bah et al. (2010). The digested peptide masses were analyzed using a MALDI-TOF-TOF mass spectrometer (ABI4700 System, USA). The corresponding parameters were based on BAH et al. (2010). Peptide mass fingerprint data were matched to the NCBIInr database using the Profound program under 50 ppm mass tolerance. Data were processed

using Data Explorer software and proteins were unambiguously identified by searching against a comprehensive non-redundant sequence database using the MASCOT software search engine (<http://www.matrixscience.com>). Proteins with change of > 1.5-fold or < -1.5-fold and a P value < 0.05 were considered differentially expressed.

Statistical analysis

Statistical analyses were performed using Data Processing System (DPS) statistical software.

RESULTS

The average number of leaf protein spots on the 2-DE gels for the Cd treatment and the control was 1594 and 1588, respectively. Of these, 94 and 64 protein spots were observed to be significantly up-regulated (>1.5-fold change) and down-regulated (<-1.5-fold change), respectively, after Cd treatment compared with the control. Fifteen (A1-A15) and 11 (B1-B11) protein spots were successfully identified by MALDI-TOF-TOF-MS (Tables 1 and 2, Figs 1 and 2). The functions of the fifteen proteins that were significantly up-regulated in the Cd treatment were categorized as metabolism (46.7%), signal transduction (26.7%), stress and defense response (6.7%), energy (6.7%) and unknown function (13.3%) (Table 1). Seven proteins were involved in metabolism: 5-methyltetrahydropteroyltriglutamate-homocysteine methyltransferase (A1), glyceraldehyde-3-phosphate dehydrogenase (A2), glutamine synthetase (A3), tubulin beta chain (A4), 2,3-bisphosphoglycerate-independent phosphoglycerate mutase (A5), alpha-1,4-glucan-protein synthase (A6) and 3 beta-hydroxysteroid-dehydrogenase/decarboxylase isoform 1 (A7). The other up-regulated proteins were as follows: signal transduction (c.f. A8 and A9, S-adenosylmethionine synthetase 1; A11, 1-aminocyclopropane-1-carboxylate oxidase), stress and defense (A12, pathogenesis-related protein 10), energy (A13, ATP synthase subunit alpha) and unknown function (A14 and A15).

Table 1. Proteins whose expression were significantly up-regulated under Cd stress

Spot ID	Protein name	Accession number	MW (Da)	pI	FC	Score	Cov %	MP	PF
A1	5-methyltetrahydropteroyltriglutamate-homocysteine methyltransferase (<i>Catharanthus roseus</i>)	METE_CATRO	85089	6.10	10 ⁶	126	11	7	M
A2	Glyceraldehyde-3-phosphate dehydrogenase, (<i>Oryza sativa</i>)	G3PC_ORYSI	36561	6.61	10 ⁶	132	10	4	M

A3	Glutamine synthetase (<i>Musa acuminata</i>)	MOS022_MUSAM	39428	6.03	1.80	94	16	4	M
A4	Tubulin beta chain (<i>Cicer arietinum</i>)	TBB_CICAR	51171	4.75	2.90	65	25	7	M
A5	2,3-bisphosphoglycerate-independent phosphoglycerate mutase (<i>Cryophytum crystallinum</i>)	PMGI_MESCR	61316	5.39	3.12	141	13	7	M
A6	Alpha-1,4-glucan-protein synthase (<i>Pisum sativum</i>)	UPTG_PEA	42059	5.73	1.50	110	17	6	M
A7	3beta-hydroxysteroid-dehydrogenase/decarboxylase isoform 1 (<i>Arabidopsis thaliana</i>)	HSDD1_ARATH	48493	8.89	3.86	29	2	1	M
A8	S-adenosylmethionine synthetase 1 (<i>Populus trichocarpa</i>)	METK1_POPTR	43642	5.68	10 ⁶	245	20	7	ST
A9	S-adenosylmethionine synthetase 1 (<i>Populus trichocarpa</i>)	METK1_POPTR	43642	5.68	4.05	236	20	7	ST
A10	Auxin-responsive protein IAA18 (<i>Arabidopsis thaliana</i>)	IAA18_ARATH	29936	9.26	2.89	61	13	6	ST
A11	1-aminocyclopropane-1-carboxylate oxidase (<i>Persea americana</i>)	ACCO_PERAE	36453	5.08	10 ⁶	42	4	1	ST
A12	Pathogenesis-related protein 10 (<i>Crocus sativus</i>)	E5D8G2_CROSA	17502	6.08	22.58	93	37	5	SD
A13	ATP synthase subunit alpha, chloroplastic (<i>Eucalyptus globulus</i>)	ATPA_EUCGG	55535	5.15	1.52	319	24	12	E
A14	Uncharacterized protein (<i>Triticum aestivum</i>)	W5A1P2_WHEAT	192765	6.74	1.80	72	12	16	U
A15	Uncharacterized protein (<i>Musa acuminata</i>)	M0TI68_MUSAM	32323	5.23	1.85	51	4	1	U

Cov, coverage; E, energy; FC, fold change; M, metabolism; MP, matched peptides; PF, Putative function; U, unknown; SD, stress and defense response; ST, signal transduction.





Figure 1. Representative 2-DE maps of saffron leaf proteins isolated from Cd and control treatment. Total proteins were extracted and separated by 2-DE. In IEF, 100 mg proteins were loaded onto pH 4–7 IPG strips (24 cm, linear). SDS-PAGE was performed with 12.5% gels. The spots were visualized by silver staining. Differentially accumulated protein spots are indicated by green sashes. Fifteen higher expressed spots (A) and eleven suppressed (B) spots are shown in the maps.

Table 2. Proteins whose expression were significantly down-regulated under Cd stress

Spot ID	Protein name	Accession number	MW (Da)	pI	FC	Score	Cov %	MP	PF
B1	Ribulose biphosphate carboxylase/oxygenase activase, (<i>Oryza sativa</i>)	RCA_ORYSJ	51764	5.43	-2.85	253	16	8	M
B2	Ribulose biphosphate carboxylase large chain (<i>Isophysis tasmanica</i>)	RBL_ISOTA	52248	5.87	-1.58	386	22	12	M
B3	Stromal 70 kDa heat shock-related protein, chloroplastic (<i>Pisum sativum</i>)	HSP7S_PEA	75583	5.22	-2.97	170	17	9	SD
B4	Eukaryotic initiation factor 4A (<i>Triticum aestivum</i>)	IF4A_WHEAT	47183	5.31	-2.11	59	14	7	PS
B5	30S ribosomal protein 2, chloroplastic (<i>Spinacia oleracea</i>)	RRP2_SPIOL	28390	8.42	-2.61	91	10	5	PS
B6	Chaperonin (<i>Zea mays</i>)	B4F848_MAIZE	25796	8.49	-2.68	76	33	6	PS
B7	Ferredoxin-NADP reductase, leaf isozyme, (<i>Pisum sativum</i>)	FENR1_PEA	40454	8.56	-7.00	118	16	7	T
B8	Ferritin-1, chloroplastic (<i>Glycine max</i>)	FRI1_SOYBN	28204	5.73	-1.50	104	9	5	T
B9	Knotted1 (<i>Spinifex sericeus</i>)	A3RCP6_9POAL	9309	4.98	-1.50	78	92	7	Tr
B10	Uncharacterized protein (<i>Hordeum vulgare</i>)	M0UGX1_HORVD	91892	6.51	-3.20	76	22	15	U
B11	Uncharacterized protein (<i>Brachypodium distachyon</i>)	I1HXX4_BRADI	39825	4.50	-2.68	72	8	5	U

Cov, coverage; FC, fold change; M, metabolism; MP, matched peptides; PF, Putative function; PS, protein synthesis; U, unknown; SD, stress and defense response; T, transport; Tr, transcription.

**Figure 2.** The spot view of individual identified spots, being up-regulated (A) and down-regulated (B) after Cd treatment.

The 11 suppressed proteins (Cd vs. control) belonged to 6 functional categories (Table 2; Fig. 2). Two were found to play key roles in photosynthesis: ribulose biphosphate carboxylase/oxygenase activase (B1) and ribulose biphosphate carboxylase large chain (B2). There were 3 protein synthesis-related proteins: eukaryotic initiation factor 4A (B4), 30S ribosomal protein 2 (B5) and chaperonin (B6). One stress and defense response protein, stromal 70 kDa heat shock-related protein (B3), and 1 transcription-related protein, Knotted1 (B9), were significantly down-regulated by -2.97-fold and -1.50-fold, respectively, after Cd treatment. Two transport-related proteins, ferredoxin-NADP reductase (B7) and ferritin-1 (B8), were also identified. In addition, 2 uncharacterized proteins (B10 and B11) were also clearly down-regulated.

DISCUSSION

Cd is believed to induce damage even at very low concentrations, not only affecting many important physiological processes but also reducing the yield and quality of plants (CAO et al., 2013, 2014). Saffron is a traditional precious herb. Our previous study found that Cd significantly suppressed saffron growth and considerable Cd was accumulated (data unpublished). In this study, 15 and 11 protein spots were up-regulated and down-regulated, respectively, in response to Cd stress (Fig. 2).

Among the 15 proteins with elevated expression, 4 are involved in signal transduction: S-adenosylmethionine synthetase 1 (SAMS, A8 & A9), 1-aminocyclopropane-1-carboxylate oxidase (ACCO, A11) and auxin-responsive protein IAA18 (A10). It has been found that the ethylene (ET) signaling pathway plays a fundamental role in the response to abiotic stress, including Cd stress (FUJITA et al. 2006). SAMS has long been considered to have a house-keeping function and plays a crucial role in ET synthesis (DAI et al. 2013). In addition, ACCO catalyzes the terminal step of ET synthesis (HERBETTE et al. 2006; CAO et al. 2014). Hence, up-regulation of SAMS and ACCO in response to Cd stress could contribute to ET synthesis, which subsequently activates additional Cd-responsive genes. Auxin-responsive protein IAA18 was also significantly up-regulated (Table 1). Therefore, it is necessary to study the relationship between the ET and auxin signaling pathways in response to Cd toxicity.

Seven of the proteins identified as up-regulated were involved in metabolism (Table 1). Glyceraldehyde-3-phosphate dehydrogenase

(GAPDH, A2) plays a key role in glycolysis. VESCOVI et al. (2013) found that the cytosolic GAPDH isoenzyme GAPC1 may play a role in signaling oxidative stress or protection in Arabidopsis roots under Cd stress. Glutamine synthetase (GS, A3) catalyzes the formation of glutamate, which is an important component of reduced glutathione and phytochelatins (PCs). Rana et al. (2008) observed that the expression of cytosolic GS was induced by Cd and salt stress. Studies have found that beta-tubulin and alpha-1,4-glucan-protein synthase are involved in plant cell wall metabolism (SPOKEVICIUS et al., 2007; JIN et al., 2011). Lee et al. (2010) also found that expression of alpha-1,4-glucan-protein synthase (A6) was significantly elevated. Membrane-bound ATPases provide energy and an H⁺ gradient for the co-transport of copper, cobalt, lead, and Cd, with protons used for the detoxification of these metal ions in plants (CAO et al. 2014). CAO et al. (2014) found that transcripts of V-ATPase were significantly elevated in a Cd-tolerant barley genotype but were not affected in a Cd-sensitive genotype. Kieffer et al. (2008) found that pathogenesis-related proteins showed a dramatic increase in expression. In our study, these proteins were all significantly up-regulated under Cd stress. Plants have evolved sophisticated defense mechanisms as an adaption to Cd stress. The results of this study suggest that Cd may induce signal molecules (ET) that subsequently activate Cd-responsive proteins, elevate the expression of antioxidant-related proteins that could scavenge the Cd-induced accumulation of ROS, and transport Cd into vacuoles and sequester Cd in cell walls, which would contribute to a decrease in the available Cd²⁺ of the cytosol.

The negative impact of Cd on photosynthesis is well documented (CLIJSTERS; VAN ASSCHE 1985). Rubisco catalyzes the first step in net photosynthetic CO₂ assimilation and photorespiratory carbon oxidation (SPREITZER and SALVUCCI 2002). Ferredoxin NADP(H) oxidoreductases (FNRs) catalyze electron transfer between NADP(H) and ferredoxin (LINTALA et al., 2007). An absence of *AtLFNR1* led to a decrease in green leaf size, lower chlorophyll content and fewer light-harvesting complex proteins, and a markedly lower PSI/PSII ratio in the mutant than the wild type (LINTALA et al., 2007). In the present study, expression of rubisco (B1, B2), ferredoxin-NADP reductase (B7) and ferritin-1 (B8) were all down-regulated after Cd treatment. These results indicate that Cd toxicity damaged the photosynthesis system in saffron. Razavizadeh et al.

(2009) found that the stromal 70 kDa heat shock protein was significantly down-regulated under salinity in tobacco leaves. The authors believed that this down-regulation was likely the cause of the failure to protect cells against salinity. Similar results were also found in this study. Heat shock protein and molecular chaperonin have been found to play vital roles in protein stabilization, folding, assembly and translocation under abiotic stress (KOMATSU; HOSSAIN, 2013). Stromal 70 kDa heat shock-related protein (B3) and chaperonin (B6) were significantly down-regulated (Table 1). The 30S ribosomal protein 2 was also down-regulated after Cd treatment. These results demonstrate that Cd stress affected the normal functioning of the

chloroplast. Moreover, down-regulation of initiation factor 4A also affected the level of translation in saffron. In summary, Cd stress affected the normal functioning of the chloroplast, reducing photosynthesis and translation.

ACKNOWLEDGEMENTS

This work was supported by the Star Program of the State Ministry of Science & Technology (2015GA700053), the Zhejiang Natural Science Foundation (LY13B020001) and the Agricultural Project of Zhejiang Public Technology Research (2015C32104).

RESUMO: O cádmio (Cd) é um poluente ambiental global altamente prejudicial. As plantas desenvolveram mecanismos de defesa complexos como uma adaptação contra a toxicidade por Cd. Neste estudo, realizou-se um experimento em vaso para avaliar o perfil proteico do açafrão em resposta ao estresse por Cd. Foi descoberto que quinze proteínas foram supra-reguladas (up-regulated) nas folhas de plantas sob estresse por Cd e foram principalmente relacionados ao metabolismo, transdução de sinal, estresse e resposta de defesa e energia. Foi descoberto ainda que onze proteínas foram infra-reguladas (down-regulated) após tratamento com Cd, incluindo ribulose bifosfato carboxilase oxigenase (RuBisCO), ferredoxina-NADP redutase, uma proteína relacionada com o choque térmico de 70 kDa e três proteínas associadas à síntese de proteínas. Os resultados fornecem informações valiosas sobre o mecanismo molecular do açafrão em resposta à toxicidade do Cd e a possível utilização de recursos genéticos no desenvolvimento de açafrão tolerante ao Cd e de baixa acumulação.

PALAVRAS-CHAVE: Metal pesado. Proteoma. Resposta. Fotossíntese. Açafrão.

REFERENCES

- ABOU-ARAB, A. A. K.; ABOU DONIA, M. A. Heavy metals in egyptian spices and medicinal plants and the effect of processing on their levels. **Journal of Agricultural and Food Chemistry** v. 48, p. 2300-2304, 2000. <http://dx.doi.org/10.1021/jf990508p>
- ASADI, M. H.; ZAFARI, F., SARVEAZAD, A.; ABBASI, M.; SAFA, M.; KORUJI, M.; YARI, A.; MIRAN, R. A. Saffron improves epididymal sperm parameters in rats exposed to cadmium. **Nephro-urology monthly**, v. 6, p. e12125, 2014. <http://dx.doi.org/10.5812/numonthly.12125>
- BAH, A. M.; SUN, H.; CHEN, F.; ZHOU, J.; DAI, H.; ZHANG, G.; WU, F. Comparative proteomic analysis of *Typha angustifolia* leaf under chromium, cadmium and lead stress. **Journal of Hazardous Materials** v. 184, p. 191-203, 2010. <http://dx.doi.org/10.1016/j.jhazmat.2010.08.023>
- CAO, F.; WANG, R.; CHENG, W.; ZENG, F.; AHMED, IM.; HU, X.; ZHANG, G.; WU, F. Genotypic and environmental variation in cadmium, chromium, lead and copper in rice and approaches for reducing the accumulation. **Science of The Total Environment** v. 496, p. 275-281, 2014a. <http://dx.doi.org/10.1016/j.scitotenv.2014.07.064>
- CAO, F.; CHEN, F.; SUN, H.; ZHANG, G.; CHEN, Z. H.; WU, F. Genome-wide transcriptome and functional analysis of two contrasting genotypes reveals key genes for cadmium tolerance in barley. **BMC Genomics**, v. 15, p. 611, 2014b. <http://dx.doi.org/10.1186/1471-2164-15-611>

- CAO, F.; WANG, N.; ZHANG, M.; DAI, H.; DAWOOD, M.; ZHANG, G.; WU, F. Comparative study of alleviating effects of GSH, Se and Zn under combined contamination of cadmium and chromium in rice (*Oryza sativa*). **Biometals**, v. 26, p. 297-308, 2013. <http://dx.doi.org/10.1007/s10534-013-9611-9>
- CARPENTIER, S. C.; WITTERS, E.; LAUKENS, K.; DECKERS, P.; SWENNEN, R.; PANIS, B. Preparation of protein extracts from recalcitrant plant tissues: An evaluation of different methods for two-dimensional gel electrophoresis analysis. **Proteomics**, v. 5, p. 2497-2507, 2005. <http://dx.doi.org/10.1002/pmic.200401222>
- CLIJSTERS, H.; Van Assche, F. Inhibition of photosynthesis by heavy metals. **Photosynthesis Research** v. 7, p. 31-40, 1985. <http://dx.doi.org/10.1007/BF00032920>
- DAI, H.; CAO, F.; CHEN, X.; ZHANG, M.; AHMED, IM.; CHEN, Z.; LI, C.; ZHANG, G.; WU, F. Comparative proteomic analysis of aluminum tolerance in tibetan wild and cultivated barleys. **PLoS One**, v. 8, p. 5, 2013. <http://dx.doi.org/10.1371/journal.pone.0063428>
- DENG, G.; Li, M.; Li, H.; Yin, L.; Li, W. Exposure to cadmium causes declines in growth and photosynthesis in the endangered aquatic fern (*Ceratopteris pteridoides*). **Aquatic Botany** v. 112, p. 23-32, 2014. <http://dx.doi.org/10.1016/j.aquabot.2013.07.003>
- FUJITA, M.; FUJITA, Y.; NOUTOSHI, Y.; TAKAHASHI, F.; NARUSAKA, Y.; YAMAGUCHI-SHINOZAKI, K.; SHINOZAKI, K. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. **Current Opinion in Plant Biology** v. 9, p. 436-442, 2006. <http://dx.doi.org/10.1016/j.pbi.2006.05.014>
- HALL, J. L. Cellular mechanisms for heavy metal detoxification and tolerance. **Journal of Experimental Botany** v. 53, p. 1-11, 2002. <http://dx.doi.org/10.1093/jxb/53.366.1>
- HERBETTE, S.; TACONNAT, L.; HUGOUVIEUX, V.; PIETTE, L.; MAGNIETTE, M. L. M.; CUINE, S.; AUROY, P.; RICHAUD, P.; FORESTIER, C.; BOURGUIGNON, J.; RENO, J. P.; VAVASSEUR, A.; LEONHARDT, N. Genome-wide transcriptome profiling of the early cadmium response of Arabidopsis roots and shoots. **Biochimie**, v. 88, p. 1751-1765, 2006. <http://dx.doi.org/10.1016/j.biochi.2006.04.018>
- JIN, H.; DO, J.; MOON, D.; NOH, E. W.; KIM, W.; KWON, M. EST analysis of functional genes associated with cell wall biosynthesis and modification in the secondary xylem of the yellow poplar (*Liriodendron tulipifera*) stem during early stage of tension wood formation. **Planta**, v. 234, p. 959-977, 2011. <http://dx.doi.org/10.1007/s00425-011-1449-1>
- KIEFFER, P.; DOMMES, J.; HOFFMANN, L.; HAUSMAN, J.; RENAUT, J. Quantitative changes in protein expression of cadmium-exposed poplar plants. **Proteomics**, v. 8, p. 2514-2530, 2008. <http://dx.doi.org/10.1002/pmic.200701110>
- KOMATSU, S.; HOSSAIN, Z. Organ-specific proteome analysis for identification of abiotic stress response mechanism in crop. **Frontiers in Plant Science** v. 4, p. 71, 2013. <http://dx.doi.org/10.3389/fpls.2013.00071>
- KULIK, A.; ANIELSKA-MAZUR, A.; BUCHOLC, M.; KOEN, E.; SZYMANSKA, K.; ZMIENKO, A.; KRZYWINSKA, E.; WAWER, I.; MCLOUGHLIN, F.; RUSZKOWSKI, D.; FIGLEROWICZ, M.; TESTERINK, C.; SKLODOWSKA, A.; WENDEHENNE, D.; DOBROWOLSKA, G. SNF1-Related protein kinases type 2 are involved in plant responses to cadmium stress. **Plant Physiology** v. 160, p. 868-883, 2012. <http://dx.doi.org/10.1104/pp.112.194472>
- KUNIHIRO, S.; SATIO, T.; MATSUDA, T.; INOUE, M.; KURAMATA, M.; TAGUCHI-SHIOBARA, F.; YOUSSEFIAN, S.; BERBERICH, T.; KUSANO, T. Rice *DEP1*, encoding a highly cysteine-rich G protein γ subunit, confers cadmium tolerance on yeast cells and plants. **Journal of Experimental Botany** v. 64, p. 4517-4527, 2013. <http://dx.doi.org/10.1093/jxb/ert267>

- LEE, K.; BAE, D. W.; KIM, S. H.; HAN, H. J.; LIU, X.; PARK, H. C.; LIM, C. O.; LEE, S. Y.; CHUNG, W. S. Comparative proteomic analysis of the short-term responses of rice roots and leaves to cadmium. **Journal of Plant Physiology** v. 167, p. 161-168, 2010. <http://dx.doi.org/10.1016/j.jplph.2009.09.006>
- LINTALA, M.; ALLAHVERDIYEVA, Y.; KIDRON, H.; PIIPPO, M.; BATTCHIKOVA, N.; SUORSA, M.; RINTAMÄKI, E.; SALMINEN, T. A.; ARO, E.; MULO, P. Structural and functional characterization of ferredoxin-NADP⁺-oxidoreductase using knock-out mutants of Arabidopsis. **The Plant Journal** v. 49, p. 1041-1052, 2007. <http://dx.doi.org/10.1111/j.1365-313X.2006.03014.x>
- MACCARONE, R.; DI MARCO, S.; BISTI, S. Saffron supplement maintains morphology and function after exposure to damaging light in mammalian retina. **Investigative Ophthalmology & Visual Science** v. 49, p. 1254-1261, 2008. <http://dx.doi.org/10.1167/iovs.07-0438>
- RANA, N. K.; MOHANPURIA, P.; YADAV, S. K. Expression of tea cytosolic glutamine synthetase is tissue specific and induced by cadmium and salt stress. **Biologia Plantarum**, v. 52, p. 361-364, 2008. <http://dx.doi.org/10.1007/s10535-008-0075-7>
- RAZAVIZADEH, R.; EHSANPOUR, A. A.; AHSAN, N.; KOMATSU, S. Proteome analysis of tobacco leaves under salt stress. **Peptides**, v. 30, p. 1651-1659, 2009. <http://dx.doi.org/10.1016/j.peptides.2009.06.023>
- SHAMSA, A.; HOSSEINZADEH, H.; MOLAEI, M.; SHAKERI, M. T.; RAJABI, O. Evaluation of *Crocus sativus* L. (saffron) on male erectile dysfunction: A pilot study. **Phytomedicine**, v. 16, p. 690-693, 2009. <http://dx.doi.org/10.1016/j.phymed.2009.03.008>
- SPOKEVICIUS, A. V.; SOUTHERTON, S. G.; MACMILLAN, C. P.; QIU, D.; GAN, S.; TIBBITS, J. F. G.; MORAN, G. F.; BOSSINGER, G. β -tubulin affects cellulose microfibril orientation in plant secondary fibre cell walls. **The Plant Journal** v. 51, p. 717-726, 2007. <http://dx.doi.org/10.1111/j.1365-313X.2007.03176.x>
- SPREITZER, R. J.; SALVUCCI, M. E. Rubisco: structure, regulatory interactions, and possibilities for a better enzyme. **Annual Review of Plant Biology** v. 53, p. 449-475, 2002. <http://dx.doi.org/10.1146/annurev.arplant.53.100301.135233>
- URAGUCHI, S.; FUJIWARA, T. Cadmium transport and tolerance in rice: perspectives for reducing grain cadmium accumulation. **Rice**, v. 5, p. 1-8, 2012. <http://dx.doi.org/10.1186/1939-8433-5-5>
- VESCOVI, M.; ZAFFAGNINI, M.; FESTA, M.; TROST, P.; LO SCHIAVO, F.; COSTA, A. Nuclear accumulation of cytosolic glyceraldehyde-3-phosphate dehydrogenase in cadmium-stressed Arabidopsis roots. **Plant Physiology**, v. 162, p. 333-346, 2013. <http://dx.doi.org/10.1104/pp.113.215194>
- VESELOV, D.; KUDOYAROVA, G.; SYMONYAN, M.; VESELOV, S. Effect of cadmium on ion uptake, transpiration and cytokinin content in wheat seedlings. **Bulgarian Journal of Plant Physiology** v. 29, p. 353-359, 2003.
- XU, C.; SIBICKY, T.; HUANG, B. Protein profile analysis of salt-responsive proteins in leaves and roots in two cultivars of creeping bentgrass differing in salinity tolerance. **Plant Cell Reports** v. 29, p. 595-615, 2010. <http://dx.doi.org/10.1007/s00299-010-0847-3>