

ANTI-INFLAMMATORY, ANTINOCICEPTIVE AND ANTIOXIDANT ACTIVITIES OF THE HYDROMETHANOLIC FRACTION FROM *ANNONA NUTANS* LEAVES

ATIVIDADE ANTI-INFLAMATÓRIA, ANTINOCICEPTIVA E ANTIOXIDANTE DA FRAÇÃO HIDROMETANÓLICA DAS FOLHAS DE *ANNONA NUTANS*

Nathalia L. SILVA¹; Aline A. SALDANHA²; Denise B. SILVA³; Carlos A. CAROLLO³; Ângela L. B. SARTORI⁴; Adriana C. SOARES⁵; João M. de SIQUEIRA⁶

1. Doutoranda, Laboratório de Farmacognosia/Química de Produtos Naturais, Campus Centro-Oeste, Universidade Federal de São João Del Rei, Divinópolis, MG, Brazil; 2. Doutoranda, Laboratório de Farmacologia da Dor e Inflamação, Campus Centro-Oeste, Universidade Federal de São João Del Rei, Divinópolis, MG, Brazil; 3. Doutor, Laboratório de Produtos Naturais e Espectrometria de Massas, Universidade Federal de Mato Grosso do Sul, Campo Grande, MS, Brazil; 4. Doutor, Centro de Ciências Biológicas e da Saúde, Universidade Federal de Mato Grosso do Sul, Campo Grande, MS, Brazil; 5. Doutor, Laboratório de Farmacologia da Dor e Inflamação, Campus Centro-Oeste, Universidade Federal de São João Del Rei, Divinópolis, MG, Brazil; 6. Doutor, Laboratório de Farmacognosia/Química de Produtos Naturais, Campus Centro-Oeste, Universidade Federal de São João Del Rei, Divinópolis, MG, Brazil. jmaximo@ufsj.edu.br

ABSTRACT: *Annona nutans* (Annonaceae) is a plant species found in Bolivia, Paraguay, Argentina, and the Brazilian Cerrado, specifically in the states of Mato Grosso and Mato Grosso do Sul (Brazil). Its common names are *Araticû-Mi* and *Araticû-Ñu*. The research contributions regarding the chemical composition and biological activities of extracts from *A. nutans* are rare, with only four articles being found in the literature. Therefore, the present study evaluated the anti-inflammatory and antinociceptive activities of the hydromethanolic fraction (FHMeOH) using carrageenan-induced paw edema and hot-plate tests. In addition, the antioxidant activity was evaluated by DPPH radical scavenging, total phenolic, flavonoid and tannin content assays and quantification of the major metabolites by LC-MS were performed. Oral treatment with the FHMeOH (at a dose of 300 mg.kg⁻¹) significantly reduced paw edema 2 h and 4 h after the inflammatory stimulus. The intraperitoneal (i.p.) treatment with the FHMeOH (50 and 100 mg.kg⁻¹) proved to be most effective, and the inhibition of acute inflammation was still visible 6 h after carrageenan injection. At doses of 50 and 100 mg.kg⁻¹ (i.p.), FHMeOH exhibits central antinociceptive effects by increasing the latency of the reaction in the hot-plate model. The FHMeOH showed antioxidant potential, and the metabolites quercetin-3-*O*-galactoside, quercetin-3-*O*-glucoside, isorhamnetin-3-*O*-galactoside, quercetin-3-*O*- β -D-apiofuranosyl-(1 \rightarrow 2)-galactopyranoside, and chlorogenic acid were identified and quantified by LC-MS. Our results indicate, for the first time, that FHMeOH has anti-inflammatory, antinociceptive and antioxidant potential, and it is a promising source of studies for new herbal medicines

KEYWORDS: *Annona nutans*. Flavonoids. Anti-inflammatory. DPPH. Antinociceptive. HPLC quantification.

INTRODUCTION

Annona is among the 27 most important genera of the Annonaceae family. This genus, containing approximately 162 species distributed among the tropics, is represented mainly in South and Central America, with 110 native tropical species in the Americas and Africa (CHATROU et al., 2012; COUVREUR et al., 2011). Initially, plants belonging the Annonaceae family were believed to predominantly contain alkaloids (LEBOEUF et al., 1982); however, a great diversity of chemical constituents was recently acknowledged. Studies of species belonging to the Annonaceae family intensified following the isolation of a chemical

group known as “annonaceous acetogenins”, which presented a wide variety of biological activities such as cytotoxic, antitumor, pesticidal, and antimicrobial (BERMEJO, 2005; CAVÉ et al., 1997). For many years, the interest in *Annona* species, and other genera in the Annonaceae family, was a result of the presence of acetogenins, which showed promising pharmacological activities (BERMEJO, 2005). However, epidemiological data from the island of Guadalupe (Caribbean) have associated the consumption of *Annona* species (*A. muricata*, *A. reticulata*, and *A. squamosa*) with the development of atypical parkinsonism, suggesting that acetogenins and quinoline alkaloid derivatives are directly related to its etiology, because the

acetogenin annonacin was shown to cause neurodegeneration in rats (CAPARROS-LEFEBVRE; STEELE, 2005). Apart from acetogenins, plants belonging to the Annonaceae family possess a wide variety of metabolites that are responsible for important pharmacological activities, such as anti-inflammatory, antinociceptive, and antioxidant (BENITES et al., 2015; FORMAGIO et al., 2013a, 2013b).

Annona nutans (or *Annona spinescens* var. *nutans*) is a plant species found in Bolivia, Paraguay, Argentina, and the Brazilian Cerrado, specifically in the states of Mato Grosso and Mato Grosso do Sul (Brazil). Its more common names are *aratico*, *chirimoya del campo*, and *sinini de la pampa*; and those in the Guarani language are *Araticû-Mi* and *Araticû-Ñu* (CORRÊA, 1926; TROPICOS.ORG, 2017). The research contributions regarding the chemical composition and biological activities of the extracts from *A. nutans* are rare, with only four articles found in the literature (GLEYE et al., 2000, 1998; GONÇALVES et al., 2014; SILVA et al., 2015). The acetogenins were found in *A. nutans* roots (GLEYE et al., 2000, 1998), with studies in the literature showing the absence of acetogenins in the leaves (SILVA et al., 2015; SILVA, 2013). Thus, the present study identified and quantified some metabolites by LC-MS and quantitated the total phenolic, tannin, and flavonoid content, in addition to assessing the antioxidant potential as DPPH radical scavenging ability from the hydromethanolic fraction (FHMeOH) of *A. nutans*. Moreover, the anti-inflammatory and antinociceptive efficacy of the FHMeOH was evaluated *in vivo*.

MATERIAL AND METHODS

Plant material

The leaves of *A. nutans* were collected in Porto Murtinho, the state of Mato Grosso do Sul, Brazil. The plant was previously identified by Renato de Mello-Silva, and a voucher specimen was deposited at the CGMS Herbarium (MS, Brazil) under number 27648. The present study obtained a Certificate of Registration from the National System for the Management of Genetic Heritage and Associated Traditional Knowledge (A90D499).

Preparation of fractions

Dried and powdered leaves (160 g) were percolated with methanol:water (9:1) (5.0 L) for 72 h at room temperature, yielding 32 g hydromethanolic extract. 28 g was dissolved in methanol:water (9:1) (500 mL), followed by

sequential partitioning with n-hexane, chloroform, and ethyl acetate (200 mL, 5x of each solvent). The final hydromethanolic fraction (FHMeOH) was dried under a vacuum in a rotatory evaporator at 45 °C, and subsequently lyophilized, yielding 21 g. For the determination of total phenolic, tannin, and flavonoid contents, a stock solution in ethanol:water (1:1) was initially prepared from the FHMeOH solution at a concentration of 5 mg.mL⁻¹, and successive dilutions at concentrations of 5.0; 2.5; 1.25; 0.625; 0.3125 mg.mL⁻¹ were prepared.

Determination of the total phenolic content

The total phenolic content was determined by spectrophotometric quantitation using the Folin-Ciocalteu reagent in 96-well microplates, as described previously (ZHANG et al., 2006).

Briefly, a 20 µL sample (prepared as previously described) and 100 µL Folin-Ciocalteu reagent (Imbralab®) were added to each well, shaken, and incubated for 5 min. Subsequently, 80 µL 7.5% Na₂CO₃ (Alphatec®) solution was added.

A calibration curve was generated using gallic acid (Cromato Produtos Químicos Ltda®) at the standard at concentrations of 1; 0.5; 0.25; 0.125; 0.0625; and 0.0312 mg.mL⁻¹. For the reagent blank, 20 µL methanol, 100 µL Folin-Ciocalteu reagent, and 80 µL 7.5% Na₂CO₃ were used. The plates were incubated in the dark at room temperature for 2 h. Readings were performed using a microplate spectrophotometer (SpectraMax®Plus384, Molecular Devices, Sunnyvale, CA, USA, Gen5 software) at $\lambda = 750$ nm. For the instrument blank, 20 µL CH₃OH:H₂O (1:1) and 180 µL distilled water were used. The total phenolic content (TPC) was determined by interpolation of the absorbance of the samples against the calibration curve obtained for the standard and are expressed as µg of gallic acid equivalents.mg⁻¹ for respective fractions. All analyses were performed in triplicate.

Determination of the total tannin content

Sample solutions were prepared at the same concentrations as described for the total phenolic quantitation; however, prior to the addition of Folin-Ciocalteu reagent, 0.01 g.mL⁻¹ hide powder (Sigma-Aldrich) was added, and the solution was shaken for 60 min on an orbital shaker. The filtrates (20 µL) were added to each well of a 96-well plate with 20 µL methanol:water (1:1) and 100 µL Folin-Ciocalteu reagent, shaken, and incubated for 5 min. Subsequently, 80 µL of 7.5% Na₂CO₃ solution was added, and the plates were incubated in the dark at room temperature for 2 h. Readings were performed using a microplate spectrophotometer at $\lambda = 750$ nm.

For the instrument blank, 20 µL of methanol:water (1:1) and 180 µL of distilled water were used.

The total tannin content (TTC) was determined from the standard curve by subtracting the calculated content of the sample solution for non-adsorbed polyphenols from the calculated TPC content (sample solution for total polyphenols). The expression is shown below. The results are expressed as µg.mL⁻¹ of sample. All analyses were performed in triplicate.

TTC = content of the sample solution for total polyphenols – content of the sample solution for non-adsorbed polyphenols in hide powder (BRASIL, 2010a; VERZA et al., 2007).

Determination of the total flavonoid content

The total flavonoid content (TFC), equivalent to quercetin, present in the fractions was determined using a previously described method (BANOV et al., 2006; BRASIL, 2010b), and expressed as µg quercetin . mg⁻¹ fraction.

For the calibration curve, quercetin (Sigma-Aldrich) was used as the standard reference flavonoid, of which a stock solution of 0.01 µg.mL⁻¹ was prepared. From this stock, dilutions were prepared at concentrations of 20, 10, 5, 4, 3, 2, 1, 0.75, 0.5 and 0.3 µg.mL⁻¹, to which 500 µL 5% (w/v) AlCl₃ solution was added. Following a 30-min incubation, the absorbance was read at 425 nm using a UV-Vis spectrophotometer (QUIMIS®) and the calibration curve was constructed.

To 100 µL 50% methanol, 100 µL FHMeOH and 500 µL 5% (w/v), AlCl₃ solution was added. Following a 30-min incubation, the absorbance was read at 425 nm using a UV-Vis spectrophotometer (QUIMIS®). All analyses were performed in triplicate.

Determination of the DPPH radical-scavenging capacity

The radical-scavenging capacity of FHMeOH was determined according to the method described by Burda and Oleszek (2001) (BURDA et al., 2001). BHT (2,6-di-tert-butyl-4-methylphenol) was used as the reference compound. FHMeOH was prepared in triplicate for each concentration (1, 10, 100, 250, and 500 µg.mL⁻¹). Each sample (75 µL) was added to three wells of a 96-well plate containing 150 µL 0.002% (w/v) DPPH-methanol solution, shaken vigorously, and incubated in the dark for 30 min. The control was prepared as above without any extract or BHT. The absorbance was measured at λ = 517 nm using a UV-Vis spectrophotometer (Biotek Power Wave XS2/US, U.S.A) and methanol was used for baseline

correction. The radical-scavenging activity is expressed as the inhibition percentage and was calculated as:

$$[1 - (\text{Abs}_{\text{control}} - \text{Abs}_{\text{sample}})] \times 100$$

where Abs_{control} = absorbance of DPPH radicals in methanol and Abs_{sample} = absorbance of fraction in methanol + DPPH. The scavenging activity is expressed as µg.mL⁻¹. IC₅₀ values (µg.mL⁻¹) were calculated using Probit analysis (FINNEY, 1980).

Identification and quantitative UFLC-DAD- ESI-QTOF-MS analysis

The identification and quantification of phytocompounds was performed on a Shimadzu Prominence UFLC TM system, equipped with a LC20AD, coupled to a diode array detector and mass spectrometer (MicroOTOF-Q III Bruker Daltonics, Billerica, USA) with an electrospray ion source. The analyses were monitored between 210 and 800 nm and the mass spectrometer operated in a negative ionization mode (*m/z* 120-1300). Chromatographic analyses were performed in a Kinetex C-18 column (Phenomenex, 2.6 µ, 150 x 2.1 mm). For quantification, FHMeOH was dissolved in methanol and water (1:1, v/v) at a concentration of 1 mg.mL⁻¹. Standard solutions at a concentration of 1 mg.mL⁻¹ were prepared with chlorogenic acid (5-*O*-caffeoylquinic acid), the flavonoids quercetin-3-*O*-galactopyranoside, quercetin-3-*O*-glucopyranoside, and isorhamnetin-3-*O*-galactopyranoside (Sigma Aldrich®). For the analytical curve, dilutions were prepared at the concentrations from 0.0488 to 200.00 µg.mL⁻¹ (0.0488, 0.0976, 0.195, 0.390, 0.781, 1.56, 3.12, 6.25, 12.50, 25.00, 50.00, 100.00, and 200.00 µg.mL⁻¹). For the isorhamnetin-3-*O*-galactopyranoside, concentrations of 12.5, 25.0, 50.0, 100.0, and 200.0 µg.mL⁻¹ were prepared. About 3 µL of sample was injected in to the column by the auto sampler. The samples were eluted through the column with a gradient mobile phase consisting of A (water 0.1% (v/v) formic acid) and B (acetonitrile: formic acid 0.1% (v/v)). The gradient elution was programmed as follows: 0–2 min B (3%); 2–25 min B (25%); 25–26 min B (80%); 26–28 min B (80%); 28–29 min B (3%); and 29–35 min B (3%). The analyses were carried out in triplicate at a flow rate of 0.3 mL.min⁻¹, at a temperature of 50 °C, with the detector set at λ = 340 nm. Calibration curves were plotted showing a linear relationship between concentrations *versus* peak areas for all reference compounds. The

attribution of the chromatographic peak was based on the retention times and confirmed by the injection of standards. The concentration of each peak was calculated from the experimental peak areas by analytical interpolation in a standard calibration line. Peak areas were calculated at 340 nm. The limit of detection (LOD) was determined as a signal-to-noise ratio of 3:1 and the limit of quantification (LOQ) was determined as a signal-to-noise ratio of 10:1 (GARCÍA-SALAS et al., 2015). The precision was calculated by relative standard deviation (%RSD), and the selectivity was evaluated by comparing the chromatograms of the individual reference standards and the degree of interference between the peaks when injected simultaneously; the degree of purity of these peaks was also investigated (BRITO, 2014; RIBANI et al., 2004).

In vivo assays

Chemicals

Indomethacin (Ind) and carrageenan λ type IV were purchased from Sigma-Aldrich Inc. (St. Louis, MO, USA). Fentanyl citrate (Fent) was purchased from Cristália (SP, Brazil). DMSO 2% in physiological saline was used as the control, and FHMeOH was prepared in this vehicle for oral or intraperitoneal treatments to mice.

Animals

Adult male Swiss mice (28–30 g) were obtained from the Bioterium of Universidade Federal de São João del-Rei, Brazil, and were housed in temperature-controlled rooms (22–25 °C), under a 12–12 h light–dark cycle, with access to food and water *ad libitum*. The mice were acclimated for one week prior to the experiment. Twelve hours prior to the beginning of oral treatments, the mice were fasted and received only water *ad libitum*. For the intraperitoneal experiment, the food and water were retained. The number of mice and the intensity of noxious stimuli used were the minimum necessary to demonstrate consistent effects of the drug treatments. All procedures were carried out in accordance with the guidelines set forth by the Brazilian National Council for the Control of Animal Experimentation and International Association for the Study of Pain and were approved by the Ethics Committee in Animal Experimentation of the Federal University of São João Del-Rei, Brazil (CEUA/UFSJ, protocol 034/2015).

Assessment of the anti-inflammatory activity

The anti-inflammatory activity was assayed

using the carrageenan-induced paw edema model (LEVY, 1969). Mice were randomly divided into five groups (n=6), and orally (p.o.) received vehicle (10 mL.kg⁻¹, control group), FHMeOH (at doses of 30, 100, and 300 mg.kg⁻¹), or indomethacin (10 mg.kg⁻¹). Moreover, FHMeOH, at doses of 25, 50 and 100 mg.kg⁻¹, were intraperitoneally (i.p.) administered. Following 30 (i.p.) or 60 (p.o.) min following treatments, carrageenan (400 µg, 30 µL) was injected into the plantar side of the left hind paw. Paw volume was measured using a plethysmometer (Insight®, Brazil) prior to treatment (basal value) and at 1, 2, 4, and 6 h after the injection of the inflammatory stimulus. The volume of edema was calculated by the difference between the prior basal paw volume and the one after carrageenan injection.

Evaluation of the antinociceptive activity

Mice were tested on a hot-plate (Insight®, Brazil) kept at a constant temperature of 55 ± 0.50 °C for 24 h before the assay, and animals that remained on the apparatus for less than 15s were selected. Thus, the selected animals were randomly divided into five groups (n=7) and received (i.p.) vehicle (10 mL.kg⁻¹, control group), FHMeOH (at doses of 25, 50, and 100 mg. kg⁻¹), and Fentanyl (Fent 200 µg.kg⁻¹). Reaction times were recorded when the mice licked their paws or jumped at intervals of 30 min up to 120 min after treatments. A cut-off of 30s was chosen to avoid tissue lesions (MUHAMMAD; SAEED; H., 2012).

Statistical analysis

Microsoft Excel 2010 (Microsoft Corporation) was used for the quantitation of the total phenolic, tannin, and flavonoid content. In the evaluation of the anti-inflammatory and antinociceptive activities, results are expressed as the mean ± SEM. The statistical significance between groups was assessed using one-way analysis of variance (ANOVA) followed by the Bonferroni multiple comparison *post-hoc* test. All calculations for anti-inflammatory, antinociceptive activities and DPPH radical-scavenging capacity were performed using GraphPad Prism™ version 5.01 (GraphPad® Software Inc., San Diego, CA). A level of significance ($p < 0.05$) was considered for each experiment.

RESULTS

Total phenolic, tannin and flavonoid content

The total phenolic, tannin, and flavonoid content of the FHMeOH extract from *A. nutans* was calculated based on the interpolation of the absorbance values of the samples from the calibration curve of gallic acid (TPC and TTC), which presented the following equation of the line, $y = 0.2352x + 0.0044$, $R^2 = 0.9982$, and from the calibration curve of quercetin (TFC), which presented $y = 0.0228x + 0.0027$, $R^2 = 0.9986$. The total phenolic, tannin, and flavonoid content was

$62.96 \pm 3.73 \mu\text{g.mg}^{-1}$, $31.14 \pm 3.11 \mu\text{g.mg}^{-1}$, and $18.07 \pm 0.10 \mu\text{g.mg}^{-1}$, respectively.

Determination of DPPH radical-scavenging capacity

The DPPH radical-scavenging activity of the FHMeOH extract from *A. nutans* is presented in Figure 1. FHMeOH showed a dose-dependent inhibitory effect with an IC_{50} of $4.89 \mu\text{g.mL}^{-1}$, which was comparable to that of the commercial antioxidant, BHT ($\text{IC}_{50} = 16.36 \pm 3.63 \mu\text{g.mL}^{-1}$). The FHMeOH at doses of 1, 10, 100 and 250 $\mu\text{g.mL}^{-1}$ presented a scavenging effect on the DPPH radical that was statistically significant compared to the BHT standard.




Figure 1. The DPPH radical-scavenging ability of the hydromethanolic fraction (FHMeOH) and 2,6-di-tert-butyl-4-methylphenol (BHT) at five different concentrations ($\mu\text{g.mL}^{-1}$). $p < 0.001$ as compared with BHT.

Identification and quantitative UFLC-DAD- ESI-QTOF-MS analysis

For the analytical curve and linearity, the regression equations were $y = 12222x - 18557$ ($R^2 = 0.9992$) for chlorogenic acid; $y = 10087x - 13571$ ($R^2 = 0.9992$) for quercetin-3-*O*-galactopyranoside; $y = 9454.4x - 13865$ ($R^2 = 0.9991$) for quercetin-3-*O*-glucoside; and $y = 8597.5x + 12942$ ($R^2 = 0.9995$) for isorhamnetin-3-*O*-galactopyranoside.

The limit of detection (LOD) for all flavonoids was $4.88 \times 10^{-2} \mu\text{g.mL}^{-1}$ and chlorogenic acid was $9.77 \times 10^{-2} \mu\text{g.mL}^{-1}$. The limit of quantification (LOQ) was $1.95 \times 10^{-1} \mu\text{g.mL}^{-1}$ for flavonoids and $3.91 \times 10^{-1} \mu\text{g.mL}^{-1}$ for chlorogenic acid. Relative standard deviations (%RSD) were in the range of 0.11% to 3.49% and they were calculated as a mean of the three replications.

The estimated concentration of metabolites was calculated in $\mu\text{g.mg}^{-1}$ of FHMeOH measuring $3.04 \mu\text{g.mg}^{-1}$ of chlorogenic acid, $6.74 \mu\text{g.mg}^{-1}$ of quercetin-3-*O*-galactopyranoside, $3.62 \mu\text{g.mg}^{-1}$ of quercetin-3-*O*-glucopyranoside, $2.12 \mu\text{g.mg}^{-1}$ of isorhamnetin-3-*O*-galactopyranoside, and 4.86

$\mu\text{g.mg}^{-1}$ of quercetin-3-*O*- β -D-apiofuranosyl-(1 \rightarrow 2)-galactopyranoside. Since the metabolite, quercetin-3-*O*- β -D-apiofuranosyl-(1 \rightarrow 2)-galactopyranoside, had no reference standard, its content was calculated based on the analytical curve of quercetin-3-*O*- β -galactopyranoside with a correction factor based on the corresponding molecular weight, because they have the same aglycone and thus the same chromophore group. This compound (quercetin-3-*O*- β -D-apiofuranosyl-(1 \rightarrow 2)-galactopyranoside) was identified in studies conducted in our laboratory using ^1H and ^{13}C NMR, COSY and DEPT techniques (SILVA et al., 2015; SILVA, 2013).

The FHMeOH was also analyzed by UFLC-DAD-MS to identify its chemical constituents. The compounds were identified by the comparison of UV spectra and retention time with applied patterns and subsequent confirmation of their molecular weights and their fragmentation in MS^2 . The compounds identified are listed in Table 1 and illustrated on the chromatogram in Figure 2.

Twenty-three compounds could be detected and identified from FHMeOH. Chlorogenic acids derivatives identified were 3-*O*-*E*-caffeoylquinic

acid (**3**) and 4-*O-E*-caffeoylquinic acid (**7**), as well as the O-glycosylated flavonols **14-22** and alkaloids **4, 6, 8-11** and **23**. These compounds were identified by comparing the spectral data reported for them in

the literature (CLIFFORD et al., 2003; LIU et al., 2018) and data were also reported from *Annona* species for the alkaloids (FERRAZ et al., 2017; SHANGGUAN et al., 2018).




Figure 2. Base peak chromatogram obtained in negative and positive ion modes from FHMeOH of *A. nutans*.

Table 1. Constituents identified by UFLC-DAD-MS from FHMeOH

Peak	RT (min)	Compound	MF	UV (nm)	Negative mode (m/z)		Positive mode (m/z)	
					MS [M-H] ⁻	MS/MS	MS [M+H] ⁺	MS/MS
1	1.2	<i>O</i> -dihexoside	C ₁₂ H ₂₂ O ₁₁	-	341.1086	191	365.1068 ^{Na}	
2	1.7	NI	C ₁₁ H ₂₂ O ₈	-	279.1087	-	281.1252	
3	6.4	3- <i>O-E</i> -caffeoylquinic acid	C ₁₆ H ₁₇ O ₉	299, 323	353.0885	191, 179	355.1043	163
4	7.5	norcoclaurine-hexoside	<i>O</i> -C ₂₇ H ₃₅ NO ₁₂	-	-	-	566.2268	272, 255, 161
5	7.9	NI	C ₁₉ H ₂₈ O ₁₁	275	431.1555	-	433.1728	205, 187, 175
6	10.1	NI	C ₁₈ H ₁₉ NO ₄	288	312.1239	-	314.1406	178, 163
7	10.7	4- <i>O-E</i> -caffeoylquinic acid st	C ₁₆ H ₁₈ O ₉	299, 325	353.0886	191, 173	355.1027	163
8	11.3	di- <i>O</i> -methoxyl hydroxyl alkaloid aporphine	di-C ₁₈ H ₁₉ NO ₄	285	312.1254	-	314.1409	298, 284, 270, 151
9	12.6	NI	C ₁₇ H ₁₉ NO ₃	285	-	-	286.1456	254, 237, 209, 191, 175, 165
10	13.0	Stepharine	C ₁₈ H ₁₉ NO ₃	285	-	-	298.1451	254, 238, 223, 161, 146
11	14.5	Magnoflorine	C ₂₀ H ₂₄ NO ₄ ⁺	280	-	-	342.1712 [*]	297, 282, 265, 237, 192
12	15.1	NI	C ₁₉ H ₂₃ NO ₄	283	328.1566	-	330.1708	284, 267, 192, 177
13	17.4	NI	C ₁₉ H ₃₂ O ₈	-	387.2021	-	389.2180	227, 209, 191, 173
14	17.5	Quercetin- <i>O</i> -hexosyl-deoxyhexoside	C ₂₇ H ₃₀ O ₁₆	270, 350	609.1443	300, 271, 255	611.1605	303
15	17.8	Quercetin- <i>O</i> -hexosyl-	C ₂₇ H ₃₀ O ₁₆	270,	609.1453	300, 271, 255	611.1623	303

		deoxyhexoside		350					
16	18.0	Quercetin-3- <i>O</i> - β -D-apiofuranosyl-(1 \rightarrow 2)-galactopyranoside	C ₂₆ H ₂₈ O ₁₆	265, 350	595.1305	300, 271, 255, 243, 179	597.1473	465, 303	
17	18.2	Quercetin- <i>O</i> -hexosyl-deoxyhexoside	C ₂₇ H ₃₀ O ₁₆	270, 350	609.1441	300, 271, 255, 179	611.1630	465, 303	
18	18.3	Quercetin- <i>O</i> -pentosyl-hexoside	C ₂₆ H ₂₈ O ₁₆	265, 350	595.1289	300, 271, 255, 243, 179	597.1485	465, 303	
19	18.5	Quercetin-3- <i>O</i> - β -galactopyranoside st	C ₂₁ H ₂₀ O ₁₂	265, 352	463.0872	300, 271, 255, 243	465.1044	303	
20	19.0	Quercetin-3- <i>O</i> - β -glucopyranoside st	C ₂₁ H ₂₀ O ₁₂	266, 348	463.0881	300, 271, 255, 243, 179	465.1044	303	
21	21.4	Quercetin- <i>O</i> -methyl- <i>O</i> -hexoside	C ₂₂ H ₂₂ O ₁₂	268, 350	477.1031	314, 299, 285, 271, 257, 243	479.1196	317, 302, 285	
22	21.8	Quercetin- <i>O</i> -methyl- <i>O</i> -hexoside	C ₂₂ H ₂₂ O ₁₂	265, 350	477.1027	314, 299, 285, 271, 257, 243	479.1201	317, 302, 285	
23	24.5	Xylopinine	C ₁₈ H ₁₇ NO ₃	284	-	-	296.1288	279, 264, 249, 234, 221, 206, 178	
25	26.2	NI	C ₅₉ H ₈₇ N ₅ O ₁₅	-	-	-	553.8154 ^{**}	482, 416, 359, 237, 209	

RT: retention time, MF: molecular formula; NI: non-identified; Na: [M+Na]⁺; *: [M]⁺; **: [M+2H]²⁺; st: confirmed by the injection of authentic standard

Anti-inflammatory effect of the FHMeOH

With respect to the anti-inflammatory activity, at 2 and 4 h post-carrageenan injection, the p.o. administration of 300 mg.kg⁻¹ FHMeOH

promoted a significant reduction in paw edema by 48.57% ($p < 0.05$) and 52.0% ($p < 0.05$), respectively, as compared with the control group (Figure 3).




Figure 3. The effect of p.o. administration of the FHMeOH from *A. nutans* on carrageenan-induced paw edema in mice. Data were analyzed by ANOVA followed by Bonferroni's multiple comparison *post-hoc* test. Values are expressed as the mean \pm SEM ($n = 6$). * $p < 0.05$ and ** $p < 0.01$ compared with the control group.

In contrast, i.p. treatment with FHMeOH at minor doses (50 and 100 mg.kg⁻¹) exerted long-lasting anti-inflammatory effects that remained significant 6 h after inflammatory stimulus (Figure

4). The inhibitory values of paw edema at 2 and 4 h post-carrageenan-induced acute inflammation were 95.45% ($p < 0.001$) and 74.29% ($p < 0.001$), respectively, for 50 mg.kg⁻¹ of the FHMeOH.




Figure 4. The effect of i.p. administration of the hydromethanolic fraction (FHMeOH) from *A. nutans* on carrageenan-induced paw edema. Data were analyzed by ANOVA followed by Bonferroni's multiple comparison *post-hoc* test. Values are expressed as the mean \pm SEM ($n = 6$). * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ as compared with the control; # $p < 0.05$ and ## $p < 0.01$ as compared with the 25 mg.kg⁻¹ FHMeOH group.

Antinociceptive activity of the FHMeOH

FHMeOH at doses of 50 and 100 mg.kg⁻¹ (i.p.) induced a significant increase in the latency of

reaction, and the central antinociceptive effects began 90 min post both treatments and were still observable after 120 min for the 50 mg.kg⁻¹ (Figure 5).




Figure 5. Effects of i.p. administration of the hydromethanolic fraction (FHMeOH) in the hot-plate model. Data were analyzed by ANOVA followed by Bonferroni's multiple comparison *post-hoc* test. Values are expressed as the mean \pm SEM ($n = 7$). * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ as compared with the control.

DISCUSSION

The UFLC analysis in FHMeOH of *A. nutans* leaves revealed that quercetin-3-*O*-galactopyranoside is the identified phenolic compound which is the major constituent. The quantification by the UFLC-DAD method was validated and showed linearity, selectivity, and precision (BRASIL, 2003; DE AMORIM et al.,

2014; LANDIM; FEITOZA; DA COSTA, 2013). From the Annonaceae family, reports including the quantitative determination of metabolites are uncommon, in particular for flavonoids (GARCÍA-SALAS et al., 2015), and most of the studies in the literature have evaluated the alkaloid and acetogenin content, because these two classes are the main ones in the species of the family (ALMEIDA, J. R. G. S.; JUNIOR, R. G. O; DE OLIVEIRA, 2015).

Currently, studies involving polyphenols such as flavonoids and chlorogenic acid derivatives have also gained great importance due to their diverse biological properties such as antioxidant, anti-inflammatory, antinociceptive, antimicrobial, and cardioprotective activity, among others (CALDERÓN-MONTAÑO et al., 2011; GALVÃO, STANLEY DE S. L. MONTEIRO et al., 2016; GARCÍA-LAFUENTE et al., 2009; GEORGIEV; ANANGA; TSOLOVA, 2014).

A direct relationship among antioxidant activity, phenolic compounds, and anti-inflammatory efficacy has been demonstrated in the literature (FORMAGIO et al., 2013a, 2013b; HIRANO et al., 2001), including in certain *Annona* species. The methanolic extract from *A. crassiflora*, for instance, has been shown to have a high total phenolic and flavonoid content (BENITES et al., 2015), and it effectively reduced paw edema and leukocyte recruitment induced by carrageenan at doses of 100 and 300 mg.kg⁻¹ (ROCHA et al., 2016). Similarly, the methanolic extract of *A. dioica* has high levels of total phenols and flavonoids, and in the concentration of 30 to 300 mg.kg⁻¹ p.o., it exhibited an anti-edematogenic effect in carrageenan-induced paw edema in a time- and dose-dependent manner (FORMAGIO et al., 2013b). *A. reticulata* has been also shown to possess strong antioxidant ability and dose-dependent inhibition of paw edema following carrageenan injection in rats (KANDIMALLA et al., 2016).

The present study demonstrated, for the first time, the antioxidant, anti-inflammatory and antinociceptive activities of the FHMeOH of *A. nutans* leaves. Moreover, the anti-inflammatory and antinociceptive activities of FHMeOH were demonstrated using acute inflammation and thermal hyperalgesia models in mice. Carrageenan injection into the paw provokes a biphasic response characterized by the initial phase (0 to 1 h) and the later phase (over 1 h). In the first phase, the release of histamine, serotonin, and bradykinins, and, to a lesser extent, prostaglandins occurs. The later phase is related to the overproduction of prostaglandins and polymorphonuclear leukocyte migration (CUZZOCREA et al., 1998). The p.o. treatment with FHMeOH failed to significantly inhibit paw edema formation with the lowest doses tested (30 and 100 mg.kg⁻¹). Only the highest dose of 300 mg/kg FHMeOH significantly inhibited paw edema 2 and 4 h after carrageenan injection. Since this inhibition was no longer observed at t = 6 h, this experiment suggests a low bioavailability by this route. On the other hand, the FHMeOH administered via i.p. exhibited significant anti-

edematogenic activity in both phases. According to the literature, in the some studies of the pharmacological effects of natural products, using these models and the intraperitoneal route, the treatments with plant extracts showed anti-inflammatory and antinociceptive activities at higher doses (75-500 mg.kg⁻¹) (ALMEIDA et al., 2012; IBRAHIM et al., 2002; NARDI et al., 2003; SADANHA et al., 2016) than those used for FHMeOH in the present study.

Regarding the antioxidant activity, the DPPH radical-scavenging potential was used and is often compared with that of butylated hydroxytoluene (BHT), a commercial antioxidant used as a food additive (BURDA et al., 2001). As for measured by DPPH radical scavenging, FHMeOH presented an IC₅₀ of 4.89 µg.mL⁻¹, which was superior to that shown by BHT (IC₅₀ = 16.36 µg.mL⁻¹). The IC₅₀ of *A. nutans* is lower than the methanolic extract (17.84 µg.mL⁻¹) of *A. dioica* leaves (FORMAGIO et al., 2013b), and *A. dioica* presented high rates of flavonoids (FORMAGIO et al., 2013b). Although flavonoids present known antioxidant activity, it seems that the diverse metabolic profile of *A. nutans* has better antioxidant activity than other *Annona* species with higher content in flavonoids. A possible cause for these data is the presence of other metabolites which presented an antioxidant action such as proaporphine alkaloids stepharine (AVULA et al., 2018; COSTA et al., 2015), aporphine magnoflorine (KUKULA-KOCH et al., 2016; NASEER et al., 2015), and oxoaporphine xylopine (COSTA et al., 2010).

As soon as we identified the anti-inflammatory activity of FHMeOH, we tested its analgesic activity, because these properties are shared by several non-steroidal anti-inflammatory drugs. The hot-plate model is a specific central antinociceptive assay, and the nociceptive response to thermal stimulus is supraspinally integrated (JULIUS, D; BASBAUM, 2001; WOOLFE; MACDNOALD, 1944). The FHMeOH produced central antinociceptive effects, verified by the increase in reaction time. However, further studies are needed to establish the possible mechanisms of the antinociceptive action of FHMeOH (SALDANHA et al., 2016).

It can be stated that inflammation and nociception are correlated, because nociception is one of the cardinal signs of inflammation (LENARDAO et al., 2016; YIMAM et al., 2016). It is also known that flavonoids and cinnamic derivatives, for example, chlorogenic acids, have anti-inflammatory and antinociceptive activities

(GARCÍA-LAFUENTE et al., 2009; GEORGIADIS et al., 2014; RATHEE et al., 2009; SERAFINI; PELUSO; RAGUZZINI, 2010; ZHAO, 2015; ZHU et al., 2013). These secondary metabolites may act via the inhibition of prostaglandin synthesis, neutrophil degranulation, and histamine, phosphodiesterase and protein kinases release (BASTOS, D. H. M.; ROGERO, M. M.; AREAS, 2009; RATHEE et al., 2009). The flavonoids and cinnamic derivatives play an important role because they present several biological actions besides anti-inflammatory and antinociceptive activities, such as antioxidant and anti-microbial effects, as well as the modulation of metabolic disorders (NAVEED et al., 2018). The antinociceptive and anti-inflammatory activities of FHMeOH can be attributed, at least in part, to the metabolites found in the fractions, such as flavonoids and chlorogenic acid derivatives.

CONCLUSION

The present study demonstrates, for the first time, that the FHMeOH fraction obtained from the leaves of *A. nutans* possesses *in vivo* anti-inflammatory and antinociceptive activities. Furthermore, the combination of phenolics present in this fraction can explain, at least partially, the effects observed. These activities raise interest in the therapeutic potential of the FHMeOH for the treatment and/or management of inflammatory and painful conditions.

ACKNOWLEDGEMENTS AND FUNDING

This work was supported by the *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior* (CAPES) in the CAPES/PNPD program, under the number 2833-2011. The authors acknowledge FAPEMIG and Federal University of São João del-Rei postgraduate fellowship and CNPq for an awarded research grant.

RESUMO: *Annona nutans* (Annonaceae) é uma espécie de planta encontrada na Bolívia, Paraguai, Argentina e no Cerrado brasileiro, especificamente nos estados de Mato Grosso e Mato Grosso do Sul (Brasil). Seus nomes mais comuns são *aratico* e *Araticû-Mi* e *Araticû-Nu*. As contribuições da pesquisa em relação à composição química e atividades biológicas dos extratos de *A. nutans* são raras, com apenas quatro artigos encontrados na literatura. Portanto, o presente estudo avaliou as atividades anti-inflamatória e antinociceptiva da fração hidrometanólica (FHMeOH) utilizando edema de pata induzido por carragenina e testes de placa quente. Além disso, a atividade antioxidante foi avaliada por meio de sequestro de radical DPPH, e foram realizados ensaios de quantificação de fenóis, flavonoides e taninos totais e quantificação dos principais metabólitos por CL-EM. O tratamento oral com a FHMeOH (na dose de 300 mg.kg⁻¹) reduziu significativamente o edema da pata 2 e 4 h após o estímulo inflamatório. Por outro lado, o tratamento intraperitoneal (i.p.) com FHMeOH (50 e 100 mg.kg⁻¹) provou ser mais eficaz e a inibição da inflamação aguda foi ainda visível 6 horas após a injeção de carragenina. Nas doses de 50 e 100 mg.kg⁻¹ (i.p.), FHMeOH exibiu efeitos antinociceptivos centrais aumentando a latência da reação no modelo de placa quente. FHMeOH apresentou potencial antioxidante e os metabólitos quercetina-3-*O*-galactosídeo, quercetina-3-*O*-glicosídeo, isoramnetina-3-*O*-galactosídeo, quercetina- 3-*O*- β -D-apiofuranosil-(1 \rightarrow 2)-galactopiranosídeo e ácido clorogênico foram identificados e quantificados por CL-EM. Nossos resultados indicam, pela primeira vez, que o FHMeOH possui efeitos anti-inflamatórios, antinociceptivos e antioxidantes, sendo uma fonte promissora de estudos para novos medicamentos fitoterápicos.

PALAVRAS-CHAVE: *Annona nutans*. Flavonóides. Anti-inflamatório. DPPH. Antinociceptiva. Quantificação por CLAE

REFERENCES

ALMEIDA, J. R. G. DA S.; ARAÚJO, E. C. DA C.; RIBEIRO, L. A. DE A.; LIMA, J. T. DE; NUNES, X. P.; CARNEIRO LÚCIO, A. S. S.; AGRA, M. DE F.; BARBOSA FILHO, J. M. Antinociceptive Activity of Ethanol Extract from *Duguetia chrysocarpa* Maas (Annonaceae). **The Scientific World Journal**, v. 2012, p. 1–6, 2012. <https://doi.org/10.1100/2012/859210>

- ALMEIDA, J. R. G. S.; JUNIOR, R. G. O; DE OLIVEIRA, A. P. **Annonaceae: Tópicos Seleccionados**. 1. ed. Curitiba: CRV, 2015.
- AMORIM, M. R. DE; RINALDO, D.; AMARAL, F. P. DO; VILEGAS, W.; MAGENTA, M. A. G.; VIEIRA, G. M.; SANTOS, L. C. DOS. HPLC-DAD based method for the quantification of flavonoids in the hydroethanolic extract of *Tonina fluviatilis* Aubl. (Eriocaulaceae) and their radical scavenging activity. **Quimica Nova**, v. 37, n. 7, p. 1122–1127, 2014.
- AVULA, B.; BAE, J. Y.; MAJRASHI, T.; WU, T. Y.; WANG, Y. H.; WANG, M.; ALI, Z.; WU, Y. C.; KHAN, I. A. Targeted and non-targeted analysis of annonaceous alkaloids and acetogenins from *Asimina* and *Annona* species using UHPLC-QToF-MS. **Journal of Pharmaceutical and Biomedical Analysis**, v. 159, p. 548–566, 2018. <https://doi.org/10.1016/j.jpba.2018.07.030>
- BANOV, D.; BABY, R.; DEL, M.; KANEKO, T. M.; VAL, M.; VELASCO, R. Caracteriza çã o do Extrato Seco de *Ginkgo biloba* L . em Formula çõ es de Uso T ó pico. v. 25, n. 2, p. 219–224, 2006.
- BASTOS, D. H. M.; ROGERO, M. M.; AREAS, J. A. G. dos alimentos no contexto de processos inflamatórios relacionados à obesidade. **Arquivos Brasileiros de Endocrinologia & Metabologia**, v. 53, n. 5, p. 646–656, 2009. <https://doi.org/10.1590/S0004-27302009000500017>
- BENITES, R.; FORMAGIO, A.; ARGANDOÑA, E.; VOLOBUFF, C.; TREVIZAN, L.; VIEIRA, M.; SILVA, M. Contents of constituents and antioxidant activity of seed and pulp extracts of *Annona coriacea* and *Annona sylvatica*. **Brazilian Journal of Biology**, v. 75, n. 3, p. 685–691, 2015. <https://doi.org/10.1590/1519-6984.21313>
- BERMEJO, A. ET AL. View Article Online / Journal Homepage / Table of Contents for this issue REVIEW Acetogenins from Annonaceae: recent progress in isolation, synthesis and mechanisms of action. **Natural Product Research**, v. 22, p. 269–303, 2005.
- BRASIL, A. N. D. V. S. Resolução - RE nº 899 - “Guia para validação de métodos analíticos e bioanalíticos”. **Ministério da Saúde**, p. 1–15, 2003.
- ____. Farmacopeia Brasileira. **Farmacopeia Brasileira, 5ª edição**, v. 2, p. 904p, 2010a.
- ____. Farmacopeia Brasileira. **Farmacopeia Brasileira, 5ª edição**, v. 1, p. 1–523, 2010b.
- BRITO, W. A. DE B. **Desenvolvimento e validação de métodos analíticos para determinação de diferentes guanilhidrazonas**. [s.l.] Universidade Federal do Rio Grande do Norte, 2014.
- BURDA, S.; BURDA, S.; OLESZEK, W.; OLESZEK, W. Antioxidant and Antiradical Activities of Flavonoids. **Journal of Agricultural and Food Chemistry**, v. 49, n. JULY 2001, p. 2774–2779, 2001. <https://doi.org/10.1021/jf001413m>
- CALDERÓN-MONTAÑO, J.; BURGOS-MORÓN, E.; PÉREZ-GUERRERO, C.; LÓPEZ-LÁZARO, M. A review on the dietary flavonoid kaempferol. **Mini reviews in medicinal chemistry**, v. 11, n. 4, p. 298–344, 2011. <https://doi.org/10.2174/138955711795305335>
- CAPARROS-LEFEBVRE, D.; STEELE, J. Atypical parkinsonism on Guadeloupe, comparison with the parkinsonism-dementia complex of Guam, and environmental toxic hypotheses. **Environmental Toxicology and Pharmacology**, v. 19, n. 3, p. 407–413, 2005. <https://doi.org/10.1016/j.etap.2004.12.052>
- CAVÉ, A.; CHABOCHE, C.; FIGADÈRE, B.; HARMANGE, J. C.; LAURENS, A.; PEYRAT, J. F.; PICHON, M.; SZLOSEK, M.; COTTE-LAFITTE, J.; QUÉRO, A. M. Study of the structure-activity relationships of the acetogenin of annonaceae, muricatacin and analogues. **European Journal of Medicinal Chemistry**, v. 32, n. 7-8, p. 617–623, 1997. [https://doi.org/10.1016/S0223-5234\(97\)83287-4](https://doi.org/10.1016/S0223-5234(97)83287-4)

- CHATROU, L. W.; PIRIE, M. D.; ERKENS, R. H. J.; COUVREUR, T. L. P.; NEUBIG, K. M.; ABBOTT, J. R.; MOLS, J. B.; MAAS, J. W.; SAUNDERS, R. M. K.; CHASE, M. W. A new subfamilial and tribal classification of the pantropical flowering plant family Annonaceae informed by molecular phylogenetics. **Botanical Journal of the Linnean Society**, v. 169, n. 1, p. 5–40, 2012. <https://doi.org/10.1111/j.1095-8339.2012.01235.x>
- CLIFFORD, M. N.; JOHNSTON, K. L.; KNIGHT, S.; KUHNERT, N. Hierarchical scheme for LC-MSⁿ identification of chlorogenic acids. **Journal of Agricultural and Food Chemistry**, v. 51, n. 10, p. 2900–2911, 2003. <https://doi.org/10.1021/jf026187q>
- CORRÊA, M. P. **Dicionário das Plantas Uteis do Brasil e das Exóticas Cultivadas**. V.I ed. Rio de Janeiro: Vol I, Ministério da Agricultura, Ed. Imprensa Nacional, 1926.
- COSTA, E. V.; PINHEIRO, M. L. B.; BARISON, A.; CAMPOS, F. R.; SALVADOR, M. J.; MAIA, B. H. L. N. S.; CABRAL, E. C.; EBERLIN, M. N. Alkaloids from the bark of *Guatteria hispida* and their evaluation as antioxidant and antimicrobial agents. **Journal of Natural Products**, v. 73, n. 6, p. 1180–1183, 2010. <https://doi.org/10.1021/np100013r>
- COSTA, E. V.; SAMPAIO, M. F. C.; SALVADOR, M. J.; NEPEL, A.; BARISON, A. Chemical constituents from the stem bark of *Annona pickelii* (Annonaceae). **Química Nova**, v. 38, n. 6, p. 769–776, 2015. <https://doi.org/10.5935/0100-4042.20150069>
- COUVREUR, T. L. P.; PIRIE, M. D.; CHATROU, L. W.; SAUNDERS, R. M. K.; SU, Y. C. F.; RICHARDSON, J. E.; ERKENS, R. H. J. Early evolutionary history of the flowering plant family Annonaceae: Steady diversification and boreotropical geodispersal. **Journal of Biogeography**, v. 38, n. 4, p. 664–680, 2011. <https://doi.org/10.1111/j.1365-2699.2010.02434.x>
- CUZZOCREA, S.; ZINGARELLI, B.; HAKE, P.; SALZMAN, A. L.; SZABO, C. Antiinflammatory effects of mercaptoethylguanidine, a combined inhibitor of nitric oxide synthase and peroxynitrite scavenger, in carrageenan-induced models of inflammation. **Free Radical Biology and Medicine**, v. 24, n. 3, p. 450–459, 1998. [https://doi.org/10.1016/S0891-5849\(97\)00280-3](https://doi.org/10.1016/S0891-5849(97)00280-3)
- FERRAZ, C. R.; SILVA, D. B.; PRADO, L. C. DA S.; CANABRAVA, H. A. N.; BISPO-DA-SILVA, L. B. Antidiarrhoeic effect and dereplication of the aqueous extract of *Annona crassiflora* (Annonaceae). **Natural Product Research**, v. 6419, n. November, p. 1–5, 2017. <https://doi.org/10.1080/14786419.2017.1396589>
- FINNEY, D. J. **Probit analysis, a statical treatment of the sigmoid response curve.**, 1980.
- FORMAGIO, A. S. N.; KASSUYA, C. A. L.; NETO, F. F.; VOLOBUFF, C. R. F.; IRIGUCHI, E. K. K.; VIEIRA, M. DO C.; FOGLIO, M. A. The flavonoid content and antiproliferative, hypoglycaemic, anti-inflammatory and free radical scavenging activities of *Annona dioica* St. Hill. **BMC complementary and alternative medicine**, v. 13, n. 1, p. 14, 2013. <https://doi.org/10.1186/1472-6882-13-14>
- FORMAGIO, A. S. N.; VIEIRA, M. DO C.; SANTOS, L. A. C. DOS; CARDOSO, C. A. L.; FOGLIO, M. A.; CARVALHO, J. E. DE; ANDRADE-SILVA, M.; KASSUYA, C. A. L. Composition and Evaluation of the Anti-Inflammatory and Anticancer Activities of the Essential Oil from *Annona sylvatica* A. St.-Hil. **Journal of Medicinal Food**, v. 16, n. 1, p. 20–25, 2013. <https://doi.org/10.1089/jmf.2011.0303>
- GALVÃO, STANLEY DE S. L. MONTEIRO, A. DE S.; SIQUEIRA, E. P.; BOMFIM, M. R. Q.; DIAS-SOUZA, M. V.; FERREIRA, G. F.; DENADAI, A. M. L.; SANTOS, ÁQUILA R. C.; SANTOS, V. L. DOS; SOUZA-FAGUNDES, E. M. DE; FERNANDES, E. S.; MONTEIRO-NETO, V. *Annona glabra* flavonoids act as antimicrobials by binding to *Pseudomonas aeruginosa* cell walls. **Frontiers in Microbiology**, v. 7, n. DEC, p. 1–9, 2016. <https://doi.org/10.3389/fmicb.2016.02053>

- GARCÍA-LAFUENTE, A.; GUILLAMÓN, E.; VILLARES, A.; ROSTAGNO, M. A.; MARTÍNEZ, J. A. Flavonoids as anti-inflammatory agents: Implications in cancer and cardiovascular disease. **Inflammation Research**, v. 58, n. 9, p. 537–552, 2009. <https://doi.org/10.1007/s00011-009-0037-3>
- GARCÍA-SALAS, P.; GÓMEZ-CARAVACA, A. M.; MORALES-SOTO, A.; SEGURA-CARRETERO, A.; FERNÁNDEZ-GUTIÉRREZ, A. Identification and quantification of phenolic and other polar compounds in the edible part of *Annona cherimola* and its by-products by HPLC-DAD-ESI-QTOF-MS. **Food Research International**, v. 78, p. 246–257, 2015. <https://doi.org/10.1016/j.foodres.2015.10.002>
- GEORGIADIS, P.; PUDNEY, P. D. A.; ROGERS, S.; THORNTON, D. J.; WAIGH, T. A. Tea derived galloylated polyphenols cross-link purified gastrointestinal mucins. **PLoS ONE**, v. 9, n. 8, 2014. <https://doi.org/10.1371/journal.pone.0105302>
- GEORGIEV, V.; ANANGA, A.; TSOLOVA, V. Recent advances and uses of grape flavonoids as nutraceuticals. **Nutrients**, v. 6, n. 1, p. 391–415, 2014. <https://doi.org/10.3390/nu6010391>
- GLEYE, C. *et al.* MURICADIENIN, MURIDIENINS AND CHATENAYTRIENINS, THE EARLY PRECURSORS OF ANNONACEOUS ACETOGENINS*. **Phytochemistry**, v. 4, n. 5, p. 749–754, 1998. [https://doi.org/10.1016/S0031-9422\(97\)00908-4](https://doi.org/10.1016/S0031-9422(97)00908-4)
- GLEYE, C.; RAYNAUD, S.; FOURNEAU, C.; LAURENS, A.; LAPRÉVOTE, O.; SERANI, L.; FOURNET, A.; HOCQUEMILLER, R. Cohibins C and D, two important metabolites in the biogenesis of acetogenins from *Annona muricata* and *Annona nutans*. **Journal of Natural Products**, v. 63, n. 9, p. 1192–1196, 2000. <https://doi.org/10.1021/np000061a>
- GONÇALVES, C. A. *et al.* Evaluation of mutagenic, teratogenic, and immunomodulatory effects of *Annona nutans* hydromethanolic fraction on pregnant mice. **Genetics and Molecular Research**, v. 13, n. 2, p. 4392–4405, 2014. <https://doi.org/10.4238/2014.June.11.3>
- GREEN, C. E.; HIBBERT, S. L.; WILLIAMS, L. A. D.; BAILEY-SHAW, Y. A.; SALMON, C.; SMITH, A. 2. Evaluation of the total phenols, total flavonoids, total tannins, and antioxidant capacity of the young vs. mature *Annona muricata* (sour sop) tea leaf grown in Jamaica. v. 661, n. 2, 2015.
- HIRANO, R.; SASAMOTO, W.; MATSUMOTO, A.; ITAKURA, H.; IGARASHI, O.; KONDO, K. Antioxidant Ability of Various Flavonoids against DPPH Radicals and LDL Oxidation. **J Nutr Sci Vitaminol**, v. 47, p. 357–362, 2001. <https://doi.org/10.3177/jnsv.47.357>
- IBRAHIM, T.; MARCOS, J.; CUNHA, T.; MADI, K.; LEA, M. B.; COSTA, S. S.; GONC, V. L. Immunomodulatory and anti-inflammatory effects of *Kalanchoe brasiliensis*. v. 2, p. 875–883, 2002. [https://doi.org/10.1016/S1567-5769\(02\)00020-6](https://doi.org/10.1016/S1567-5769(02)00020-6)
- JULIUS, D.; BASBAUM, A. I. Molecular mechanisms of nociception. **Nature**, v. 413, n. September, p. 203–210, 2001. <https://doi.org/10.1038/35093019>
- KANDIMALLA, R.; DASH, S.; KALITA, S.; CHOUDHURY, B.; MALAMPATI, S.; KALITA, K.; KOTOKY, J. Bioactive guided fractions of *Annona reticulata* L. bark: Protection against liver toxicity and inflammation through inhibiting oxidative stress and proinflammatory cytokines. **Frontiers in Pharmacology**, v. 7, n. JUN, p. 1–12, 2016. <https://doi.org/10.3389/fphar.2016.00168>
- KUKULA-KOCH, W.; KOCH, W.; ANGELIS, A.; HALABALAKI, M.; ALIGIANNIS, N. Application of pH-zone refining hydrostatic countercurrent chromatography (hCCC) for the recovery of antioxidant phenolics and the isolation of alkaloids from *Siberian barberry* herb. **Food Chemistry**, v. 203, p. 394–401, 2016. <https://doi.org/10.1016/j.foodchem.2016.02.096>

- LANDIM, L. P.; FEITOZA, G. S.; COSTA, J. G. M. DA. Development and validation of a HPLC method for the quantification of three flavonoids in a crude extract of *Dimorphandra gardneriana*. **Revista Brasileira de Farmacognosia**, v. 23, n. 1, p. 58–64, 2013. <https://doi.org/10.1590/S0102-695X2012005000111>
- LEBOEUF, M.; CAVE, A.; BHAUMIK, P. K.; MUKHERJEE, B.; MUKHERJEE, R. The phytochemistry of the annonaceae. **Phytochemistry**, v. 21, n. 12, p. 2783–2813, 1982. [https://doi.org/10.1016/0031-9422\(80\)85046-1](https://doi.org/10.1016/0031-9422(80)85046-1)
- LENARDAO, E. J.; SAVEGNAGO, L.; JACOB, R. G.; VICTORIA, F. N. Antinociceptive Effect of Essential Oils and Their Constituents: an Update Review. **Journal of the Brazilian Chemical Society**, v. 27, n. 3, p. 435–474, 2016. <https://doi.org/10.5935/0103-5053.20150332>
- LEVY, L. Carrageenan paw edema in the mouse. **Life Sciences**, v. 8, n. 11, p. 601–606, 1969. [https://doi.org/10.1016/0024-3205\(69\)90021-6](https://doi.org/10.1016/0024-3205(69)90021-6)
- LIU, M.; HUANG, X.; LIU, Q.; CHEN, M.; LIAO, S.; ZHU, F.; SHI, S.; YANG, H.; CHEN, X. Rapid screening and identification of antioxidants in the leaves of *Malus hupehensis* using off-line two-dimensional HPLC–UV–MS/MS coupled with a 1,1'-diphenyl-2-picrylhydrazyl assay. **Journal of Separation Science**, v. 41, n. 12, p. 2536–2543, 2018. <https://doi.org/10.1002/jssc.201800007>
- MORI, L. S.; BOLLER, S.; KASSUYA, C. A. L.; STEFANELLO, M. E. A.; ZAMPRONIO, A. R. Phytomedicine Analgesic effects of the ethanolic extract from *Magnolia ovata* (Magnoliaceae) trunk bark and of N-acetylxylopine , a semi-synthetic analogue of xylopine. **Phytomedicine**, v. 18, p. 143–147, 2011. <https://doi.org/10.1016/j.phymed.2010.06.001>
- MUHAMMAD, N.; SAEED, M.; H., K. Antipyretic, analgesic and anti-inflammatory activity of *Viola betonicifolia* whole plant. **BMC complementary and alternative medicine**, v. 12, p. 1–8, 2012. <https://doi.org/10.1186/1472-6882-12-59>
- NARDI, G.; FELIPPI, R.; DALBO, S.; SIQUEIRA-JUNIOR, J.; ARRUDA, D. C.; DELLE MONACHE, F.; TIMBOLA, A. K.; PIZZOLATTI, M. G.; CKLESS, K.; RIBEIRO-DO-VALLE, R. M. Anti-inflammatory and antioxidant effects of *Croton celtidifolius* bark. **Phytomedicine**, v. 10, p. 176–184, 2003. <https://doi.org/10.1078/094471103321659906>
- NASEER, S.; LONE, S. H.; LONE, J. A.; KHUROO, M. A.; BHAT, K. A. LC-MS guided isolation, quantification and antioxidant evaluation of bioactive principles from *Epimedium elatum*. **Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences**, v. 989, n. 1553, p. 62–70, 2015. <https://doi.org/10.1016/j.jchromb.2015.02.046>
- NAVEED, M. *et al.* Chlorogenic acid (CGA): A pharmacological review and call for further research. **Biomedicine and Pharmacotherapy**, v. 97, n. August 2017, p. 67–74, 2018. <https://doi.org/10.1016/j.biopha.2017.10.064>
- RATHEE, P.; CHAUDHARY, H.; RATHEE, S.; RATHEE, D.; KUMAR, V.; KOHLI, K. Mechanism of Action of Flavonoids as Anti-inflammatory Agents: A Review. **Inflammation & Allergy - Drug Targets**, v. 8, n. 3, p. 229–235, 2009. <https://doi.org/10.2174/187152809788681029>
- RIBANI, M.; GRESPAN BOTTOLI, C. B.; COLLINS, C. H.; FONTES JARDIM, I. C. S.; COSTA MELO, L. F. Validação em métodos cromatográficos e eletroforéticos. **Química Nova**, v. 27, n. 5, p. 771–780, 2004. <https://doi.org/10.1590/S0100-40422004000500017>

ROCHA, R. S.; KASSUYA, C. A. L.; FORMAGIO, A. S. N.; MAURO, M. D. O.; ANDRADE-SILVA, M.; MONREAL, A. C. D.; CUNHA-LAURA, A. L.; VIEIRA, M. D. C.; OLIVEIRA, R. J. Analysis of the anti-inflammatory and chemopreventive potential and description of the antimutagenic mode of action of the *Annona crassiflora* methanolic extract. **Pharmaceutical Biology**, v. 54, n. 1, p. 35–47, 2016. <https://doi.org/10.3109/13880209.2015.1014567>

SADANHA, A. A. *et al.* Chemical composition and anti-inflammatory activity of the leaves of *Byrsonima verbascifolia*. **Journal natural medicine**, 2016.

SALDANHA, A. A.; SIQUEIRA, J. M.; CASTRO, A. H. F.; MATOS, N. A.; KLEIN, A.; SILVA, D. B.; CAROLLO, C. A.; SOARES, A. C. Peripheral and central antinociceptive effects of the butanolic fraction of *Byrsonima verbascifolia* leaves on nociception-induced models in mice. **Inflammopharmacology**, 2016. <https://doi.org/10.1007/s10787-016-0300-5>

SERAFINI, M.; PELUSO, I.; RAGUZZINI, A. Flavonoids as anti-inflammatory agents. **Proceedings of the Nutrition Society**, v. 69, n. 03, p. 273–278, 2010. <https://doi.org/10.1017/S002966511000162X>

SHANGGUAN, Y.; HE, J.; KANG, Y.; WANG, Y.; YANG, P.; GUO, J.; HUANG, J. Structural Characterisation of Alkaloids in Leaves and Roots of *Stephania kwangsiensis* by LC-QTOF-MS. **Phytochemical Analysis**, v. 29, n. 1, p. 101–111, 2018. <https://doi.org/10.1002/pca.2718>

SILVA, N. L. **Estudo fitoquímico das folhas de *Annona nutans* (R . E . Fries)**. [s.l.] Universidade Federal de São João Del-Rei, 2013.

SILVA, N. L. ; ZOBIOLE, N. N. ; SILVA, D. B. ; SARTORI, A. L. . B. ; OLIVEIRA, J. ; PINTO, M. E. A. . ; SANTOS, F. J. L. . DOS; SIQUEIRA, J. M. DE. CONSTITUINTES QUÍMICOS E ATIVIDADE FITOTÓXICA DAS FOLHAS DE *Annona nutans*. v. 38, n. 3, p. 370–377, 2015.

TITOVA, M. V; RESHETNYAK, O. V; OSIPOVA, E. A.; SHUMILO, N. A.; ORESHNIKOV, A. V; NOSOV, A. M. Submerged Cultivation of *Stephania glabra* (Roxb .) Miers Cells in Different Systems : Specific Features of Growth and Accumulation of Alkaloid Stepharine. **Applied Biochemistry and Microbiology**, v. 48, n. 7, p. 645–649, 2012. <https://doi.org/10.1134/S0003683812070046>

Tropicos.org. Disponível em: <www.tropicos.org>. Acesso em: 1 ago. 2017.

VERZA, S. G.; KREINECKER, M. T.; REIS, V.; HENRIQUES, A. T.; ORTEGA, G. G. Avaliação das variáveis analíticas do método de folin-ciocalteu para determinação do teor de taninos totais utilizando como modelo o extrato aquoso de folhas de *Psidium guajava* L. **Quimica Nova**, v. 30, n. 4, p. 815–820, 2007. <https://doi.org/10.1590/S0100-40422007000400011>

WOOLFE, G.; MACDNOALD, A. D. The evaluation of the analgesic action of pethidine hydrochloride (demerol). **The Journal of Pharmacology and Experimental Therapeutics**, v. 80, n. 3, p. 300–307, 1944.

YIMAM, M.; LEE, Y.-C.; JIAO, P.; HONG, M.; NAM, J.-B.; BROWNELL, L.; HYUN, E.; JIA, Q. UP1306, a botanical composition with analgesic and anti-inflammatory effect. **Pharmacognosy Research**, v. 8, n. 3, p. 186, 2016. <https://doi.org/10.4103/0974-8490.182918>

ZHANG, Q.; ZHANG, J.; SHEN, J.; SILVA, A.; DENNIS, D. A.; BARROW, C. J. A simple 96-well microplate method for estimation of total polyphenol content in seaweeds. **Journal of Applied Phycology**, v. 18, n. 3-5, p. 445–450, 2006. <https://doi.org/10.1007/s10811-006-9048-4>

ZHAO, J. Flavonoid transport mechanisms: How to go, and with whom. **Trends in Plant Science**, v. 20, n. 9, p. 576–585, 2015. <https://doi.org/10.1016/j.tplants.2015.06.007>

ZHU, B.; ED, P.; KUMAR, S.; PANDEY, A. K. Flavonoids : Reviews Chemistry and Biological Activities of Flavonoids : An Overview. v. 2013, n. 3, p. 519–534, 2013.