

Epidemiology of traumatic spinal injuries in Brazil: systematic review

Ricardo Vieira Botelho¹, Luciana Dini Gianini Albuquerque²,
Rafael Bastianello Junior², Aluizio Augusto Arantes Júnior³

ABSTRACT

Prevention remains the most effective way to reduce the burden of spinal injuries (SI). Estimating the incidence and causes of SI is essential for the development of prevention programs. To reveal the current state of Brazilian SI epidemiology. To evaluate the current state of Brazilian SI epidemiology, we performed an electronic search using the Lilacs database (Latin American and Caribbean Health Sciences Literature). Fifteen papers and one unsubmitted ongoing study (BH project) reported the epidemiologic status of SI in Brazil. The mean age of SI patients was 34,75 years, and 84% of patients were male. The most commonly affected segment was the cervical spine (36.65%). The incidence ranged from 16 to 26 patients per million per year (average 21 per million per year). The average rate of complete spinal cord injury among five papers was 34%. The average mortality in four papers was 11.58%. Analyzing published national epidemiological data enables the estimation of the current state of Brazilian SI epidemiology.

KEYWORDS

Epidemiology, spinal cord injuries, spinal fractures.

RESUMO

Epidemiologia do trauma raquimedular no Brasil: revisão sistemática

A prevenção continua a ser a forma mais eficaz de reduzir o ônus das lesões na coluna vertebral (TRM). A estimativa da incidência e das causas do TRM é essencial para o desenvolvimento de programas de prevenção. Para revelar os dados epidemiológicos do TRM no Brasil, foi realizada uma busca eletrônica utilizando o banco de dados Lilacs (Literatura do Caribe em Ciências da Saúde da América Latina e Caribe). Quinze artigos publicados em um estudo em andamento (projeto BH) relataram a situação epidemiológica do TRM no Brasil. A idade média dos pacientes foi de 34,75 anos; 84% dos pacientes eram do sexo masculino. O segmento vertebral mais afetado foi a coluna cervical (36,65%). A incidência variou de 16 a 26 pacientes por milhão por ano (média de 21 por milhão por ano). A média de lesão medular completa entre cinco trabalhos que descreveram a variável foi de 34%. A média de mortalidade em quatro trabalhos foi de 11,58%. A análise dos trabalhos permite estimar os dados epidemiológicos nacionais, revelando sua distribuição e gravidade, possibilitando comparações com estudos futuros.

PALAVRAS-CHAVE

Epidemiologia, traumatismos da medula espinal, fraturas de coluna vertebral.

- 1 Professor, permanent advisor, Post-Graduation Program in Health Sciences – Instituto de Assistência Médica ao Servidor Público Estadual (Iamspe), São Paulo, SP, Brazil.
- 2 Fellow, Programa Institucional de Bolsas de Iniciação Científica/Conselho Nacional de Desenvolvimento Científico e Tecnológico (Pibic/CNPq), Post-Graduation Program in Health Sciences – Iamspe, São Paulo, SP, Brazil.
- 3 Professor, Medicine School, Minas Gerais Federal University (UFMG), Belo Horizonte, MG, Brazil.

Introduction

Traumatic spinal and spinal cord injuries (SI) are important causes of morbidity and mortality and a major cause of loss of independence and self-esteem in young adults.¹ Prevention remains the most effective way to reduce the SI burden. The causes of trauma depend on the types of housing present in large urban centers as well as the population's habits and education level.¹ Estimating the incidence and causes of SI is essential for the development of prevention programs.² No recent studies have detailed SI epidemiology across Brazil. However, it is possible to pool the data from several published reports from different cities to estimate the country's epidemiological status.¹⁻¹⁵ The objective of this study was to reveal the current state of Brazilian SI epidemiology.

Methods

To evaluate Brazilian SI epidemiology, we performed an electronic search using the Lilacs database (Latin American and Caribbean Health Sciences Literature), between January 1964 to October 2012. The search strategy involving MeSH terms and Portuguese keywords was as follows: "*Traumatismos da medula espinal*" ("spinal cord injuries") AND "*estudos de incidência*" ("incidence studies") OR "*estudos de prevalência*" ("prevalence studies") OR "*epidemiologia*" ("epidemiology") OR "*epidemiologia da fratura da coluna espinal*" ("spine fracture epidemiology") OR "*trauma da coluna*" ("spinal trauma").

Data from ongoing unpublished epidemiological studies were included with the permission of the authors. Studies that described series of cases from isolated institutions that were not of a strictly epidemiological nature were excluded.

Results

Fifteen papers and one unsubmitted ongoing study in the city of Belo Horizonte (BH project) yielded data on the epidemiologic status of SI in Brazil. The majority of the papers were based on regional hospital registry data. One paper was based on the responses to a questionnaire administered to all hospitals in the National Public Health system.² Another was a cross-sectional study detailing the prevalence of hospitalized SCI patients.³ One paper specifically described the

epidemiology related to diving in shallow water in the city of Ribeirão Preto.⁴ Another study described the epidemiology of SI by fire arm projectiles (FAP) in a hospital in Curitiba.⁵ Data from these works, due to their specificity, were not pooled together with the other papers.

Studied regions

Geographically and economically, Brazil is divided into five regions: southeast, south, middle west, north and northeast. We identified at least one SI study from each region (Figure 1).

Southeast region: six studies reported SI epidemiology data from the southeast region, five in the state of São Paulo,^{1,6-9} and one (BH project) in the state of Minas Gerais, in the capital, Belo Horizonte.*

Middle west region: one study described SI epidemiology in the state of Goiás, city of Goiânia.¹⁰

North region: Two papers were published about the same city (Belém) in the state of Pará.¹¹ Another study was performed in São Luís, in the state of Maranhão.

Northeast region: three papers were performed in the northeastern states of Piauí,¹² Pernambuco¹³ and Sergipe.¹⁴

Nationally based papers: two papers were based on national information. One was based on patients with spinal cord injuries who were treated in the hospitals belonging to the National Public Health System (SUS),² whereas the other was based on SI rehabilitation hospitals throughout the country.³

Gender and age

Data from five studies and the BH project^{1-3,10,13} revealed a mean age of 34,75 years for SI patients. Eighty-four percent of patients were male.

Incidence of spinal injuries

In three hospital-based studies and the BH project, the population of the city was provided, permitting estimates of SI incidence.^{1,6,12} In Piauí, the incidence was 16 SI cases per million inhabitants per year. In the region of Francisco Morato, the incidence was approximately 20 cases per million inhabitants, and in the northern city of São Paulo, the incidence of serious cervical SI was approximately 11 per million per year. Because cervical traumas comprised half of all injuries discussed in that study, the estimated incidence was 22 per million per year. The BH project counted 224 injured patients

* Data from this study were kindly supplied by Dr. Aluizio Arantes (personal communication).

over eight months. Eighteen percent (18.3%) of these patients were from Belo Horizonte. The city's estimated population in 2010 was 2,238,526 citizens, making the annual incidence 26 per million per year (25.57).

In these studies, the incidence ranged from 16 to 26 patients per million per year, with an average of 21 per million per year. The national studies^{2,3} were considered cross-sectional studies of admitted patients with spinal cord injuries.

Causes

In nine studies (Recife, Goiânia, Belém, northern São Paulo, Francisco Morato, southern São Paulo, of São Paulo, Santo André, São Luís and Teresina), the most frequent cause of injury was a fall from a height, mainly from rooftops (Table 1). This is a unique consequence

of the habitation types in the suburbs of large Brazilian urban centers. The rooftop is often compatible with leisure activities. As a result, falls from rooftops are relatively frequent. For this reason, rooftop falls have become a national public health problem. The rates and distributions of SI causes are described in table 1 and figure 1.

In Belo Horizonte and Aracaju, the main cause of SI was car accidents (39.8% and 40.8%, respectively). In one of the national studies,³ the main cause was motor vehicle accidents (41.7%).

Injuries by firearm projectile (FAP) caused 4.76% of traumas in Recife, 13% in Goiás, 36.7% in the central city of São Paulo (4) (1990), 8.1% in Heliópolis (south part of São Paulo), 18% in Santo André, 11.9% in Teresina and 27.5% in Sergipe. In Belém (2002), 13% of traumas were described as caused by violence at work. In the national study by da Paz *et al.*,³ 26.9% were caused by FAP.




Figure 1 – Regional distribution of Spinal trauma among Brazilian states/cities.

Table 1 – Causes of SI injuries organized by states/cities in studies based on direct hospital database samples

State/City	Falls	Auto	Motorcycle	Gunshot	Diving	Other
Pernambuco/Recife	1.91	4.76	19.05	4.76	–	9.5
Goiás/Goiânia	31	44	17	13	9	8
São Paulo/São Paulo (Downtown)	14.8			26.9	9.3	–
Pará/Belém (2002)	41.2	23.1		23.7	5	13
Pará/Belém (2003)	39.6	8.4		24.4	7.12	3.6
São Paulo/Francisco Morato	45.1	18.7				
São Paulo/São Paulo (South)	63	10	14	18	–	–
Piauí/Teresina	39.6	24.4		24.4	7.12	3.6
São Paulo/Santo André	47	14	10	18	–	–
São Paulo/São Paulo (North)	39	20	4	2.3	10	24.7
Sergipe/Aracaju	44	18.5	11.2	16.3	24.5	15.2
Minas Gerais/Belo Horizonte	34.8	39.8	–	9.5	2.9	6.6*
Maranhão/São Luís	42.6	24.2	17.2	12.6	3.4	

* Unpublished epidemiological study.

Affected spine segments

The most commonly affected spinal segment was the cervical spine (Table 2). In ten studies, injuries in the cervical spine occurred at a mean rate of 36.65% (ranging from 26.4 to 62.9%) (Table 2).

Table 2 – Percentage of cervical spine lesions by city

City	% patients with lesions in the cervical spine
Aracaju	48.3
Belo Horizonte	62.9*
Teresina	51
Santo André (spinal cord injuries)	50
Belém (2003)	30.4
Belém (2002)	36.25
Goiânia	45
Francisco Morato	36.1*
São Paulo (South)	39.5
São Luís	26.4

* Pooled cervical and cervico-thoracic injuries.

Complete SI rates

The percentage of patients with complete neurological lesions (tetraplegia or paraplegia, with complete sensitivity deficits) are presented for each city (Table 3). The studies carried out in Recife, Goiânia, Francisco Morato, Teresina and Belo Horizonte described the percentage of patients with complete spinal cord injuries. Data based on spinal cord injury papers were excluded, as were data from

Recife¹³ (this paper analyzed only thoracic and lumbar lesions, with 2.3% being complete lesions) and data from the north zone of the city of São Paulo¹ (this paper analyzed only cervical lesions). The average rate of complete spinal cord injury in the remaining five papers was 34%.

Table 3 – Percentage of complete spinal cord injuries

City	% ASIA A
Goiânia	43
Belém (2002)	26
Francisco Morato	26.1
Teresina	50
Belo Horizonte	23
Mean	34

Mortality rates

Three papers and the BH project reported their in-hospital mortality rates. Barros *et al.*,⁷ Santos *et al.*¹ and Pereira *et al.*¹³ reported rates of 21%, 15.2% and 10%, respectively. The BH study reported a mortality rate of 13.8%. The average mortality among the four papers was 11.58%.

Epidemiology in the State of São Paulo

The greatest number of studies (five) were performed in the southeastern state of São Paulo.^{1,6-9} Three of these studies were based on hospitals located in three different regions of the capital, São Paulo.^{1,7,8} Two other papers were conducted in the metropolitan area around São Paulo city.^{6,9}